

Evaluation of CSR Breeds for their Biology, Performance and Economic Importance of Silkworm, *Bombyx mori* L.

Murali S^{1*}, Tayal MK², Anil Dhar², Arti Devi³ and Sardar Singh²

¹Scientist - B, Regional Sericultural Research Station, Central Silk Board, Jammu, Jammu and Kashmir, India

²Scientist - D, Regional Sericultural Research Station, Central Silk Board, Jammu, Jammu and Kashmir, India

³Rearing Assistant, Regional Sericultural Research Station, Central Silk Board, Jammu, Jammu and Kashmir, India

*Corresponding Author: Murali S, Regional Sericultural Research Station, Central Silk Board, Jammu and Kashmir, India.

E-mail: dr.mmrl@rediffmail.com

Received: December 21, 2017; Published: February 12, 2018

Abstract

The classification and characterization of silkworm breeds are important for sericulture, which is supported by the constant development of new hybrids. In this study, 5 parental CSR breeds of *Bombyx mori* L. from the germplasm bank of RSRS, Miran Sahib were evaluated regarding phenotypic and biological/economic parameters of CSR breeds. Observations on the different phenotypic and economic traits of silkworm, *Bombyx mori* L. were taken. The perusal of the data reveals that the fecundity was recorded from 365.00 (CSR - 18) to 460.67 (CSR - 5) and hatching per cent ranged from 93.87 (CSR 19) to 95.88 (CSR 2). The larval weight varied in the range of 34.94 to 40.86 g whereas, larval duration was observed in the range of 24.00 days to 26.22 days. The larval length recorded as on 6th day 7.76 cm (CSR - 19) to 8.26 cm (CSR - 2). Single cocoon weight 1.49 to 1.70 g. The highest single cocoon weight was CSR - 18 (1.70 g), Single shell weight 0.28 to 0.36 g. Maximum shell weight recorded in CSR - 18 (0.36 g), Shell percentage 18.86 per cent (CSR - 4) to 21.29 per cent (CSR - 19). Maximum yield CSR - 2 (15.23 Kg), was observed significantly superior compared to others. These findings will help the sericulturists in finding suitable breed for getting more economic returns from silkworm rearing. It is observed that CSR - 2 is having highest yield per 10, 000 larvae with shell ratio 20.06 (%).

Keywords: Fecundity; Larval Duration; Larval Length; Cocoon Yield; Shell Ratio; *Bombyx mori* L.

Highlights

Five CSR breeds of *B. mori* has been evaluated and each one showed significant difference among them. However, the CSR- 2 revealed highest highest yield per 10, 000 larvae with shell ratio 20.06 (%).

Introduction

India is the second largest producer of mulberry raw silk in the country and largest silk consumers; it is also the world's largest importers of silk. As per the statistics available with Central silk board, more than 90% of the silk produced in the country is of multivoltines x bivoltine hybrid cocoons. The silk produced from such cocoons falling into the lower grades in the international grades for silk and cannot be suitable to use in the power looms especially for warp threads and they prefer imported silk for want of quality. Estimated demand of international quality silk in our country is around 6000 mt. It is a well-known fact that bivoltine silk excels in quality and productivity; it is inevitable to go in for large-scale production of bivoltine sericulture in the country. Elite bivoltine silkworm breeds are the basic and important input for the production of high-grade raw silk. Keeping in view of the demand for quality silk in the country an attempt made by Basavaraja, *et al.* [1]; Datta, *et al.* [2,3]; Datta, *et al.* [4,5]; Suresh Kumar, *et al.* [6,7]; Mal Reddy, *et al.* [8]; Dandin, *et al.* [9] led to the evolution of highly productive CSR bivoltine breeds which have potential to produce international grade silk.

Silkworm is the larvae of insects, which belongs to the order Lepidoptera experiencing complete metamorphosis. The mulberry silkworm *Bombyx mori* L. (Lepidoptera: Bombycidae) is a monophagous insect that feeds exclusively on mulberry foliage for its nutrition and produces the naturally proteinous silk. Nutritional intake has direct impact on the overall genetic traits such as larval and cocoon weight, amount of silk production, pupation, and reproductive traits. Silkworms go through four stages of development- eggs, larva, pupa and adult. The silkworm larval life cycle is divided into five instars, separated by four molts. Four distinct stages of development completes one generation of the species. The total silkworm life cycle is completed in 56 - 62 days.

An extensive study is needed to improve existing strains for commercial purposes and to develop new strains through breeding programs aimed at improving silk productivity, adaptability to local environments, and disease resistance/tolerance capabilities [10,11]. Several studies related to the use of productivity markers [12,13] morphological dissimilarity [14-17] as indicators of the best strains for breeding.

Material and Methods

The investigation was carried out on five *B. mori* parental CSR breeds viz., CSR - 2, CSR - 4, CSR - 5, CSR - 18, CSR - 19 maintained at the RSRS, Jammu during spring, 2016 and were incubated for 9 - 12 days in a neat and clean, disinfected room at 80 - 85% humidity

and 24 - 25°C temperature with 18 hrs light till pin head stage, at this stage black - boxing was done to ensure maximum hatching on exposure to bright light. The hatched larvae were reared separately under uniform laboratory conditions as described by Yokoyama [18] and Krishnaswami [19]. During the entire period of research, same micro-climate and feeding conditions were ensured as per the larval stage. The experiment analysed for phenotypic characteristics of egg, larvae and cocoon patterns (Table 1) and biological studies of larvae includes feeding duration (Table 2), hatching percentage, larval weight (Table 5) and total larval duration of different instars were studied, which reflects their variation among the breeds (Table 3) and cocoon characteristics of all breeds were recorded.

All the breeds were reared in three replications by following standard rearing techniques [19]. Three hundred larvae were retained after 2nd moult in each replication. The data pertaining to the morphology/phenotypic and biological parameters were recorded. During the entire period of research, same micro-climate and feeding conditions were ensured as per the larval stage.

At egg stage: Egg shape, egg colour, hatching percentage and average fecundity per female moth were studied.

At larval stage: Larval colour, markings, larval length and mean weight of 10 larvae on each day of V instar were studied and analyzed for different races.

The statistical analysis was done with the help of software SPSS and weight of larvae was measured with electronic balance.

At cocoon stage: Cocoon shape, cocoon colour, cocoon grain and economic characters of cocoon were noted.

Results and Discussion

Phenotypic Qualitative Parameters

- **Egg:** All the CSR breeds which are selected are ellipsoidal in shape, grey in color and shell color was recorded as white in CSR - 2, CSR - 4, CSR - 5 whereas CSR - 18 and CSR - 19 yellow in color (Table 1). Similar results were obtained by Anita., *et al* [20].
- **Larvae:** Color of newly hatched larvae was recorded as black and haemolymph color was transparent in all the breeds. The pattern of larvae was observed plain in CSR - 2, CSR - 4, CSR-5 and others it was recorded as plain-marked i.e. CSR - 18 and CSR - 19 (Table 1). The results were agreement with work done by Anita., *et al* [20].
- **Cocoon:** All the CSR breeds were distinctly white in color in CSR - 2, CSR - 4, CSR - 5 and others it was recorded as creamish i.e. CSR - 18 and CSR - 19 respectively. The shape of cocoon was dumbell and constricted (DC) in CSR - 4, CSR - 5, CSR - 19 and oval in CSR - 2 and CSR -18 respectively. The build and grain of among all the breeds was recorded hard and medium respectively (Table 1). Similar results were obtained by Anita., *et al* [20].

Breeds	Egg			Larvae			Cocoon			
	Shape	Shell Colour	Egg colour	Colour of Newly Hatched	Haemoly-mph Colour	Larval pattern	Colour	Shape	Build	Grains
CSR -2	E	W	G	B	T	P	W	O	H	M
CSR -4	E	W	G	B	T	P	W	DC	H	M
CSR -5	E	W	G	B	T	P	W	DC	H	M
CSR -18	E	Y	G	B	T	PM	C	O	H	M
CSR -19	E	Y	G	B	T	PM	C	DC	H	M

Table 1: Phenotypic qualitative characteristics of egg; larvae and cocoon parameters of CSR breeds.

Note: E: Ellipsoid; W: White; Y: Yellow; G: Grey; SS: Sand Stone; B: Black; T- Transparent; O: Oval; M: Medium; DC: Dumbell Constricted; H: Hard; P: Plain; M: Marked; C: Creamish

Biological parameters of larvae

Feeding duration

- **First instar:** The active period of feeding among all the breeds was recorded 3 days 5 h (Table 2).
- **Second instar:** 2 days 14 h were recorded as most active period of feeding among all the breeds (Table 2).
- **Third instar:** The active feeding period was recorded among all breeds was 3 days 16 h except in CSR - 5 i.e. 3 days 11 h respectively (Table 2).
- **Fourth instar:** The feeding period was observed among all breeds was varied 4 days 8 h was observed in CSR - 2, CSR - 4, CSR - 18 and CSR - 5 (4:41 h) and CSR - 19 (4:10 h) respectively (Table 2).
- **Fifth instar:** There is variation among breeds was recorded, 8 days 22 h more active feeding period was recorded in CSR- 2 and less i.e. 6 days was observed in CSR - 4 respectively (Table 2).
- The total average feeding duration among all the breeds was

more in CSR - 19 i.e. 21 days 22 h and less was recorded in CSR - 4 i.e. 19 days 19 h respectively (Table 2).

Larval duration

- **First instar:** The period among all the breeds was recorded as 4 days 5 h respectively (Table 3).
- **Second instar:** Three days 14 h were recorded as 2nd stage larval period among all the breeds (Table 3).
- **Third instar:** 4 days 16 h were recorded as 3rd stage larval period among all the breeds except in CSR - 5 i.e. 4 days 12 h (Table 3).
- **Fourth instar:** The larval period was observed 5 days 13 h among all the breeds except in CSR - 5 i.e. 6 days 18 h (Table 3).
- **Fifth instar:** There is variation among breeds, 7 days was recorded more in CSR - 5, CSR - 18, CSR - 19 and less i.e. 6 days was observed in CSR - 4 respectively (Table 3).

Race/Breed	Different stages (Days: h ± SD)					Total Avg. duration (Days: h)
	1 st instar	2 nd instar	3 rd instar	4 th instar	5 th instar	
CSR -2	3:05 ± 0.00	2:14 ± 0.00	3:16 ± 0.00	4:08 ± 0.00	8:22 ± 0.52	20:14 ± 0.08
CSR -4	3:05 ± 0.00	2:14 ± 0.00	3:16 ± 0.00	4:08 ± 0.00	6:00 ± 0.00	19:19 ± 0.00
CSR -5	3:05 ± 0.00	2:14 ± 0.00	3:11 ± 0.09	4:41 ± 0.52	7:00 ± 0.00	21:16 ± 0.52
CSR -18	3:05 ± 0.00	2:14 ± 0.00	3:16 ± 0.00	4:08 ± 0.00	7:00 ± 0.00	20:19 ± 0.00
CSR -19	3:05 ± 0.00	2:14 ± 0.00	3:16 ± 0.00	4:10 ± 0.03	7:00 ± 0.00	21:22 ± 0.47

Table 2: Data showing feeding duration among different stages of CSR breeds during spring rearing (2016).

Races/breeds	Different stages (Days: h ± SD)					Total duration (Days : h)
	1 st instar	2 nd instar	3 rd instar	4 th instar	5 th instar	
CSR -2	4:05 ± 0.00	3:14 ± 0.00	4:16 ± 0.00	5:13 ± 0.00	6:70 ± 0.52	24:70 ± 0.52
CSR -4	4:05 ± 0.00	3:14 ± 0.00	4:16 ± 0.00	5:13 ± 0.00	6:00 ± 0.00	24:00 ± 0.00
CSR -5	4:05 ± 0.00	3:14 ± 0.00	4:12 ± 0.07	6:18 ± 0.51	7:00 ± 0.00	25:00 ± 0.00
CSR -18	4:05 ± 0.00	3:14 ± 0.00	4:16 ± 0.00	5:13 ± 0.00	7:00 ± 0.00	25:00 ± 0.00
CSR -19	4:05 ± 0.00	3:14 ± 0.00	4:16 ± 0.00	5:13 ± 0.00	7:00 ± 0.00	25:00 ± 0.00

Table 3: Data showing larval duration among different stages of CSR breeds during spring rearing (2016).

Larval duration of total average among all the breeds was more in CSR - 2 i.e. 26 days twenty 2 h and less was recorded in CSR - 4 i.e. 24 days respectively (Table 5) and it was ranges from 24.00 - 26.22 days. The results were agreement with Krishnaswami [19] reported that the larval duration was longer in race M-5 (24.17), but it was non-agreement with results reported by Bothikar, *et al.* [21] the larvae reared on S -1635 recorded 19.66 days. Under ideal conditions it has been reported that the total larval duration is 25 - 30 days for selected JAM breeds (Raina, 2000). The results were justified with work done by Anita, *et al.* [20] reported similar results on JAM breeds, the larval duration which ranges between 23.22 - 26.16 days.

Larval length

The length of 5th instar larvae was recorded from first day to sixth day results revealed that there is significant differences were observed among all the breeds except in first day (Table 4). During first day the larval length ranges from 4.28 (CSR - 18) to 4.38

cm (CSR - 2), 2nd day it was ranges from 4.38 cm (CSR - 5) to 5.22 cm (CSR - 19), 3rd day ranges between 6.06 (CSR - 18) to 6.50 cm (CSR - 19), 4th day recorded as 6.40 cm (CSR - 5) to 7.42 cm (CSR - 2), 5th day it was ranges from 7.16 (CSR - 19) to 7.68 cm (CSR - 2) and 6th day recorded as 7.76 cm (CSR - 19) to 8.26 cm (CSR - 2) (Table 4). Similar results were recorded by 6.71 to 7.25 cm (Prabu, *et al.* 2011) and 6.12 to 7.05 cm [22]. The data shows that there is increase in larval length from day 1 to day 6 of 5th instar. The length of silkworm depends on amount of food it consumes. Length of larvae increases until larvae reaches its spinning state (6th day). When it reaches spinning stage the larvae reduce in size to one third of its normal length, it is the characteristic feature of a silkworm. This reduction in size increases pressure on silk glands to eject silk from the glands. Hence there is a sudden decrease in length on 6th day onwards. This decrease in length continues until pupal stage. The results also agreement with work done by Venugopal Reddy, *et al.* [23] revealed that the larval length varies from 5.00 to 7.43 cm.

Breeds	Days (5 th instar) (Length - Cm ± SD)					
	1	2	3	4	5	6
CSR -2	4.38 ± 0.08	4.60 ± 0.30	6.28 ± 0.15	7.42 ± 0.08	7.68 ± 0.08	8.26 ± 0.11
CSR -4	4.32 ± 0.13	4.56 ± 0.27	6.08 ± 0.24	7.06 ± 0.21	7.46 ± 0.11	7.98 ± 0.16
CSR -5	4.36 ± 0.13	4.38 ± 0.16	6.38 ± 0.19	6.40 ± 0.12	7.32 ± 0.15	7.78 ± 0.13
CSR -18	4.28 ± 0.08	5.20 ± 0.22	6.06 ± 0.21	6.72 ± 0.13	7.24 ± 0.11	7.80 ± 0.10
CSR -19	4.34 ± 0.05	5.22 ± 0.16	6.50 ± 0.16	6.44 ± 0.11	7.16 ± 0.17	7.76 ± 0.11
C.D. @ 1%	-	0.30	0.25	0.18	0.17	0.16
SE. m ±	0.04	0.10	0.08	0.06	0.05	0.05
C.V. (%)	2.35	4.82	3.06	2.02	1.74	1.59

Table 4: Data showing length of fifth instar larvae of CSR breeds during spring (2016).

Economic parameters

Nine economic traits of data pertaining to evaluation viz., hatching %, larval duration (h), larval weight (g), cocoon yield per 10,000 larvae by number, cocoon yield per 10,000 larvae by weight, pupation rate, cocoon weight, cocoon shell weight, and cocoon shell ratio of five breeds were presented in table 5. The perusal of the data reveals that the fecundity was recorded from 365.00 (CSR - 18) to 473.00 (CSR - 5) which shows statistically significant among all the breeds and hatching per cent ranged from 93.87 (CSR - 19) to 95.88 (CSR - 2) and showing statistically non-significant among all the breeds whereas larval duration shows statistically significant among all the breeds and recorded to a maximum of 26.22 (CSR - 2) and minimum of 24.00 (CSR - 4) whereas larval weight was recorded to a maximum of 40.86 g (CSR - 2) and minimum of 34.94 g (CSR - 5) showing statistically significant among all the breeds whereas pupation rate showing statistically non-significant among all the breeds and it was recorded as highest 91 per cent (CSR - 18) and less 89.00 per cent (CSR - 2)

and pupal weight was ranges between 12.66g (CSR - 19) to 13.46g (CSR - 18) showing statistically non-significant among all the breeds. With regard to yield per 10,000 larvae by number, CSR - 18 was recorded the highest (9333.33) and lowest in CSR - 4 (9053.33). Yield per 10,000 larvae by weight (Kg), ranged to the maximum of 15.23 Kg in CSR - 2 and minimum of 10.88 Kg in CSR - 5 showing statistically significant among all the breeds with regard both by number and weight basis. The weight of cocoon found to the highest of 17.33g in CSR - 18 and lowest of 16.05g in CSR - 4 shows statistically non-significant among all the breeds. The cocoon weight ranged from the maximum of 1.70 g (CSR - 18) and minimum of 1.49 g (CSR - 4). The shell weight was maximum (0.36 g) in CSR - 18 and minimum in CSR - 4 (0.28 g) shows statistically significant among all the breeds with respect to both cocoon weight and shell weight. The shell ratio (%) shows statistically significant among all the breeds and was highest in CSR - 19 (21.29) and lowest in CSR - 4 (18.86) respectively.

Breeds	Fecundity by No.	Hatching (%)	Larval weight (g)	Larval duration (Days : h)	Pupal weight (g)	Pupation rate (%)	Yield/ 10000 Larvae		Weight of cocoon (g)	Single cocoon weight (g)	Single shell weight (g)	Shell ratio (%)
							No.	Wt. (Kg)				
CSR 2	424.33 (20.61)	95.88 (78.31)	40.86	24.70	12.85	89.00 (70.61)	9226.67 (96.06)	15.23	16.48	1.65	0.33	20.06 (26.58)
CSR 4	443.33 (21.07)	95.49 (77.70)	35.36	24.00	13.31	89.67 (71.25)	9053.33 (95.15)	12.36	16.05	1.49	0.28	18.86 (25.72)
CSR 5	473.00 (21.77)	95.28 (77.43)	34.94	25.00	13.29	90.33 (71.86)	9173.33 (95.78)	10.88	16.61	1.65	0.32	19.40 (26.11)
CSR 18	365.00 (19.12)	94.65 (76.66)	38.36	25.00	13.46	91.00 (72.53)	9333.33 (96.61)	14.37	17.33	1.70	0.36	20.97 (27.24)
CSR 19	440.00 (20.99)	93.87 (75.63)	39.48	25.00	12.66	90.33 (71.86)	9200.00 (95.92)	14.31	16.09	1.57	0.33	21.29 (27.46)
C.D. @ 1%	0.85	-	2.36	0.42	-	-	0.83	2.01	-	0.12	0.02	1.01
SE. m±	0.26	0.60	0.73	0.13	0.47	0.54	0.26	0.63	0.52	0.04	0.007	0.31
C.V. (%)	2.23	1.36	3.38	0.93	6.26	1.32	0.47	8.14	5.52	4.27	3.81	2.06

Table 5: Economic parameters of CSR breeds reared during spring (2016).

Average fecundity per moth was highest 522 in Jam 2 and lowest was 355 in Jam 18 reported by Anita., *et al.* [20] these results were non-agreement with our results. The lowest hatching percent (93.79%) was obtained in Jam 11 race and the highest (94.17%) was obtained in Jam 27 race, with a mean of 93.92 per cent in all studied races Anita., *et al.* [20]. It was justified with our results, 93.87 (CSR - 19) to 95.88 (CSR - 2).

The larval weight was non-agreement with results reported by Bothikar., *et al.* [21] i.e. 40.54 g which were reared on S - 1635 and other breeds which on par with silkworm reared on variety M - 5 and breeds having larval weight ranges between 34.94 g (CSR - 5) to 40.86 g (CSR - 2). Similar results were recorded by Pakhale., *et al.* [24], the larval weight was ranges from 33.77 - 40.67 g.

The larval duration was non-agreement with results reported by Pakhale., *et al.* [24], the larval duration ranges from 21.04 - 22.28 days, our results shows that 24.00 - 26.22 days among the breeds.

The single cocoon weight was not justified with results reported by Bothikar., *et al.* [21] i.e. 1.86 g which were reared on S - 1635 and Rayer [25] and Chakravorty [26] reported the highest single cocoon weight on variety V- 1 and also not justified with results reported by Pakhale., *et al.* [24] single cocoon weight ranges from 1.76 - 1.86 g.

The single shell weight was non-agreement with Bothikar., *et al.* [21] report the silkworms which reared on S-1635 but it was similar results with variety M - 5 reported by Bothikar., *et al.* [21]. Rayer [25] and Chakravorty [26] reported the highest single shell weight on variety V- 1 and also not justified with results reported by Pakhale., *et al.* [24] single cocoon weight ranges from 0.32 - 0.36 g.

The cocoon shell percentage varies between 19.93 to 17.84 per cent reported by Bothikar, *et al.* [21] and it was agreement with our results the shell percentage varies between 18.86 - 21.29 per cent. Rayer [25] and Chakravorty [26] reported the cocoon shell percentage.

Cocoon yield per 10,000 larvae brushed varied in the range of 18.66 kg to 16.59 kg reported by Bothikar, *et al.* [21] and it was non-agreement with our results on S- 1635 but similar results which reared on variety M - 5, the yield varies between 9.97- 15.95 kg. Similar results reported by Rayer [25] and Chakravorty [26] reported the cocoon shell percentage.

Varietal differences for studied traits in *B. mori* has been reported by Ahsan, *et al.* [27], Li, *et al.* [11]; Furdai, *et al.* [28]. Similar studies on varietal diversity have also been sustained by the findings of Reza, *et al.* [29], Mistri and Jayaswal [30]; Ahsan, *et al.* [31]; Umashankara and Subramanya [32]; Nezhad, *et al.* 2009; Nguku, *et al.* [33]; Nguku, *et al.* [34]; Zannata, *et al.* [35] and Pal and Moorthy [36].

Conclusion

The obtained data showed that there are highly significant differences among the breeds for all the studied characters. There is a high positive correlation between economic parameters among all the breeds studied. The differences in obtained results are due to the variability and genotype characters for each individual of every breed.

Acknowledgements

The authors wish to thank the authorities of Regional Sericultural Research Station, CSB, Miran Sahib, Jammu - 181101, J and K, India for their encouragement to undertake this work. The authors are also thankful to Dr. Anil Dhar, Scientist-D, for his assistance in conducting effective work and close monitoring during the study. We acknowledge all technical staff who assisted their help in rearing.

Bibliography

- Basavaraja HK, *et al.* "New productive bivoltine hybrids". *Indian Silk* 34 (1995): 5-9.
- Datta RK, *et al.* "Evolution of new productive bivoltine hybrids, CSR2 x CSR4 and CSR2 x CSR5". *Sericologia* 40 (2000a): 151-167.
- Datta RK, *et al.* "Evolution of new productive bivoltine hybrid CSR3 x CSR6". *Sericologia* 40 (2000b): 407-416.
- Datta RK, *et al.* "Breeding of new productive hybrid CSR12 x CSR6 of silkworm *Bombyx mori* L". *International Journal of Industrial Entomology* 3 (2001a): 127-133.
- Datta RK, *et al.* "CSR18 x CSR19 - A robust bivoltine hybrid for all season rearing in the tropics". *Indian Silk* 39 (2001b): 5-7.
- Suresh Kumar N, *et al.* "On the breeding of CSR18 x CSR19 - A robust bivoltine hybrid silkworm, *Bombyx mori* L. for the tropics". *International Journal of Industrial Entomology* 5 (2002): 155-162.
- Suresh Kumar N, *et al.* "Development of new robust bivoltine hybrid (CSR46 x CSR47) of *Bombyx mori* L. for the tropics". *Indian Journal of Sericulture* 45.1 (2006): 21-29.
- Mal Reddy N, *et al.* "Breeding of productive bivoltine hybrid, CSR16 x CSR17 of silkworm, *Bombyx mori* L". *International Journal of Industrial Entomology* 8 (2003): 129-133.
- Dandin SB, *et al.* "Development of new silkworm hybrid, Chamaraja (CSR50 x CSR51) of *Bombyx mori* L. for tropics". *Indian Journal of Sericulture* 45.1 (2006): 35-44.
- Sen R, *et al.* "Induction of resistance to *Bombyx mori* nuclear polyhedrosis virus, into a susceptible bivoltine silkworm breed". *Indian Journal of Sericulture* 38 (1999): 107-112.
- Li M, *et al.* "Studies on some special characters in the silkworm (*Bombyx mori* L.) germplasm in China". *Sericologia* 41 (2001): 527-35.
- Fragoso DB, *et al.* "Selecao na evolucao de Resistencia a organofosforados em leucoptera coffeella (Guerin-Meneville) (Lepidoptera: Lyonetiidae)". *Neotropical Entomology* 32.2 (2003): 329-334.
- Marco Jr P, *et al.* "Thermoregulatory constraints on behavior: patterns in a neotropical dragonfly assemblage". *Neotropical Entomology* 34.2 (2005): 155-162.
- Aagaard K, *et al.* "Phylogenetic relationships in brown argus butterflies (Lepidoptera: Lycaenidae: Aricia) from North Western Europe". *Biological Journal of the Linnean Society* 75.1 (2002): 27-37.
- Pilgrim EM, *et al.* "Combining DNA sequences and morphology in systematics: testing the validity of the dragonfly species, *Cordulegaster bilineata*". *Heredity* 89.3 (2002): 184-190.
- Dujardin J and Le Pont F. "Geographic variation of metric properties within the neotropical sandflies". *Infection, Genetics and Evolution* 4.4 (2004): 353-359.
- Arruda VM, *et al.* "Analise morfologica da veneno de *Apis mellifera* L. (Hymenoptera: Apidae) em populacoes de Mato Grosso do Sul". *Neotropical Entomology* 36 (2007): 203-209.

18. Yokayama. Sericulture Annual Review. Entomology (1963): 287-298.
19. Krishnaswami S. "New technology of silkworm rearing". Bulletin of the Central Sericulture Research and Training Institute, Mysore (1978): 1-10.
20. Anita Tripathi NK, *et al.* "Comparative analysis of developmental stages in six different bivoltine races of *Bombyx mori*". *International Journal of Recent Scientific Research* 5.11 (2014): 1969-1972.
21. Bothikar PA, *et al.* "Growth and development of silk worm (*Bombyx mori* L.) on mulberry leaves". *Journal of Industrial Pollution Control* 30.2 (2014): 239-241.
22. Balasundaram D, *et al.* "Studies on the nutritional supplementation of vitamin C treated MR2 mulberry leaves fed by V instar larvae of silkworm, *Bombyx mori* (L.) (Lepidoptera: Bombycidae) in relation to feed efficacy and growth rate". *International Journal of Research in Biotechnology and Biochemistry* 3.1 (2013): 11-18.
23. Venugopal Reddy B, *et al.* "Quantitative profile Analysis of Mulberry Silkworm, *Bombyx mori* L. (CSR2XCSR4)". *International Letters of Natural Science* 34 (2015): 34-41.
24. Pakhale SG, *et al.* "Evaluation of Some Mulberry Varieties for Rearing Performance and Economic Traits of Silkworm (*Bombyx Mori* L.)". *Journal of Environmental Science, Toxicology and Food Technology (IOSR-JESTFT)* 8.12 (2014): 63-65.
25. Rayer SG. "Stability of improved mulberry varieties to shootlet feeding and its impact on cocoon production". *Journal of Maharashtra Agricultural Universities* 32.3 (2006): 367-369.
26. Chakravorty NJ. "Studies on different parameters of *B. mori*". *Seridoc* 20.2 (2004): 21-23.
27. Ahsan MK, *et al.* "Variability of some quantitative traits in the hybrids of silkworm, *Bombyx mori* L". *University Journal of Zoology, Rajshahi University* 19 (2000): 20-24.
28. Furdui EM, *et al.* "Comparative study of biological characters of larvae, crude and dried cocoon in 7 races of Silkworm *Bombyx mori* L. raised in Transylvania area". *Scientific Papers: Animal Science and Biotechnologies* 43.1 (2010): 490-493.
29. Reza AMS, *et al.* "Studies on the variation of some larval traits in different breeds of silkworm, *Bombyx mori* L". *University Journal of Zoology, Rajshahi University* 12 (1993): 21-24.
30. Mistri PK and Jayaswal KP. "Studies on phenotypic correlations between some economic traits of silkworm, *Bombyx mori* L". *Bulletin Series Research* 3 (1992): 26-29.
31. Ahsan MK, *et al.* "Inheritance of some quantitative traits in fifteen indigenous varieties of silkworm, *Bombyx mori* L". *University Journal of Zoology, Rajshahi University* 18 (1999): 79-83.
32. Umashankara ML and Subramanya G. "Correlation between larval weight and cocoon characters in five breeds of silkworm, *Bombyx mori* L". *Geobios* 29 (2002): 154-157.
33. Nguku EK, *et al.* "Larvae, cocoon and post-cocoon Characteristics of *Bombyx mori* L. (Lepidoptera:bombyxidae) fed on mulberry leaves fortified with Kenyan royal jelly". *Journal of Applied Sciences and Environmental Management* 11.4 (2007): 85-89.
34. Nguku EK, *et al.* "Performance of six Bivoltine *Bombyx mori*. (Lepidoptera: Bombyxidae) silkworm strains in Kenya". *The Open Entomology Journal* 3 (2009): 1-6.
35. Zanatta DB, *et al.* "Evaluation of economically important traits from sixteen parental strains of the silkworm, *Bombyx mori* L. (Lepidoptera: Bombyxidae)". *Neotropical Entomology* 38.3 (2009): 327-331.
36. Pal NB and Moorthy SM. "Assessment of variability in larval and cocoon traits in some genotypes of bivoltine silkworm, *Bombyx mori* L". *International Journal of Research in Biological Sciences* 1.4 (2011): 59-65.

Volume 2 Issue 3 March 2018

© All rights are reserved by Murali S., *et al.*