



Stress-Adaptive Microbes for Plant Growth Promotion and Alleviation of Drought Stress in Plants

Ajar Nath Yadav^{1*} and Neelam Yadav²

¹Department of Biotechnology, Akal College of Agriculture, Eternal University, Baru Sahib, Sirmour, Himachal Pradesh, India

²Gopi Nath P.G. College, Deoli-Salamatpur, Ghazipur, Uttar Pradesh, India

*Corresponding Author: Ajar Nath Yadav, Microbial Biotechnology Laboratory, Department of Biotechnology, Akal College of Agriculture, Eternal University, Baru Sahib, India.

Received: April 26, 2018; Published: May 28, 2018

Abstract

Drought is one of the major abiotic stresses accepted as the main constraint for loss of the crop yield worldwide. Further, problems are created by nutrient limitations and particularly low phosphorus (P) soil status due to which modern agricultural systems are highly dependent on chemical fertilizers. The biotechnology offers a number of sustainable solutions to mitigate these problems by using plant growth promoting (PGP) microbes. These microbes help the crops to tolerate drought conditions by different mechanisms including the production of the exopolysaccharides (EPS), phytohormones (Auxin, Gibberellic acid and Cytokinin), 1-aminocyclopropane-1-carboxylate (ACC) deaminase, solubilization of phosphorus, potassium and zinc, biological nitrogen fixation and enhancement of nutrient uptake, induction of the accumulation of osmolytes, antioxidants, upregulation or down regulation of the stress responsive genes. Inoculating plants with PGP microbes can increase tolerance against abiotic stresses such as drought, salinity and metal toxicity. The PGP microbes play important role in plant growth and soil health, which belong to diverse genera *Arthrobacter*, *Azospirillum*, *Azotobacter*, *Bacillus*, *Burkholderia*, *Enterobacter*, *Klebsiella*, *Lysinibacillus*, *Paenibacillus*, *Pseudomonas*, *Serratia*, and *Streptomyces*.

Keywords: Alleviation; Diversity; Drought Stress; Plant Growth Promotion; Stress-Adaptive Microbes

Introduction

There are number factors leading to the food crisis all over the world such as increasing urbanization and industrialization has led to the shrinkage of the land in past few decades. Increasing human population has further added to the world's food security concern and these factors are affecting the climate in a severe way. Additionally, the use of the chemical fertilizers is also affecting the environment. The pressure of population is increasing with such an alarming rate that there is a critical need to enhance the agricultural productivity in an eco-friendly manner. Agriculture is one of the most exposed sectors to various climatic changes. One of the major hurdle to increase the yield and productivity is exposure of the crops to the drought conditions in different parts of the world [1-5]. The drought is one of the major abiotic stresses acting as the limiting factor affecting the agricultural productivity worldwide. It has been estimated that there is approximately 9 - 10% reduction in the national production of the cereals due to the drought conditions [6].

Plant-associated beneficial microbes are recently attaining greater attention as they play an important role in enhancing the productivity of the crops and also providing resistance against the stress conditions and are known as plant growth promoting microbes (PGPMs) [7,8]. The PGPMs contribute to mitigate the stress conditions by diverse mechanisms [9,10]. The PGPMs directly enhancing the uptake of the micronutrients, through phytohormones production; fixing of atmospheric nitrogen; P, K, and Zn-solubilization or indirectly stimulating the immune system against various fungal pathogens by production of various compounds, enzymes, siderophores, antibiotics, osmolytes or improving either texture or structure of the soil [11].

Phosphorus (P) is the major macronutrient which is required by the plants for their various metabolic processes including energy transfer, signal transduction, macro-molecular biosynthesis, photosynthesis and respiration but is simultaneously the major limiting mineral nutrient for the growth of the plants due to its least availability as well as the least mobility. The replacement of soil P reserves through chemical fertilization is common, but long-term practice. There are many studies which have reported that beneficial microbes are efficient in solubilizing nutrients from soil [12-19]. The solubilization of inorganic insoluble phosphate salts by microbes result in the production or release of organic acid and organic acid decreases the pH [20-22]. The major P-solubilizers belong to genera *Achromobacter*, *Acinetobacter*, *Agrobacterium*, *Arthrobacter*, *Aspergillus*, *Azospirillum*, *Azotobacter*, *Bacillus*, *Enterobacter*, *Erwinia*, *Flavobacterium*, *Haloarcula*, *Halobacterium*, *Halococcus*, *Micrococcus*, *Mycobacterium*, *Penicillium*, *Pseudomonas*, *Rhizobium* and *Serratia* [21,23-27,28-31]. Various mechanisms used by P-solubilizers to convert the insoluble forms of the phosphorus into the soluble forms consist of acidification, chelation, exchange reactions and production of organic acids [22,32].

Another important role of plant growth promoting rhizobacteria is the synthesis of various phytohormones (plant growth regulators). The well-known phytohormones includes auxins most common being the indole acetic acid, cytokinins, gibberellins. There are diverse bacterial genera such as *Acinetobacter*, *Arthrobacter*, *Bacillus*, *Corynebacterium*, *Delftia*, *Duganella*, *Exiguobacterium*, *Kocuria*, *Lysinibacillus*, *Methylobacterium*, *Micrococcus*, *Micrococcus*, *Paenibacillus*, *Paenibacillus*, *Pantoea*, *Pseudomonas*, *Psychrobacter*, *Serratia* and *Stenotrophomonas* has been reported to produced diverse group of phytohormones [33-40]. Iron is one

of the most vital elements important for the growth of all living organisms. It acts as the cofactor for different enzymes, it is involved in process of photosynthesis, respiration, nitrogen fixation and its deficiency leads to various metabolic alterations [41]. Iron is present in abundance in the soil but is not available for the plants as well as the microbes present in the soil as the oxidized form of the iron which is Fe^{3+} reacts forming oxides and hydroxides which is not accessible to the plants as well as the microbes. Under such iron limiting conditions PGPR have the capacity to produce low molecular weight iron chelating compounds called as the siderophores for the acquisition of the ferric ions [42].

Microbe-mediated drought stress

The elucidation of the various mechanisms by which plants respond to drought stress is very important that stress tolerant plants could be grown. This process is very complex as it involves various factors which are affecting and at the same time the factors which are affected. During drought the availability of the nutrients is also affected, and this can be overcome by the use of the plant growth promoting microbes [43-45]. Ethylene when present at high concentration proves to be inhibitory for the growth of the plants. But, PGP microbes possess an enzyme 1-aminocyclopropane-1-carboxylate (ACC) deaminase enzyme which converts ACC, the immediate precursor of ethylene to α -ketobutyrate and ammonium thus lowering the concentration of the ethylene during the stress conditions and stimulating the growth of the plants. ACC deaminase activity has been reported in *Achromobacter xylosoxidans*, *Agrobacterium genomovars*, *Alcaligenes*, *Azospirillum lipoferum*, *Bacillus licheniformis*, *Brachybacterium saurashtrense*, *Brevibacterium casei*, *Brevibacterium iodinum*, *Burkholderia phytofirmans*, *Cronobacter sakazakii*, *Enterobacter cloacae*, *Methylobacterium fujisawaense*, *Pseudomonas putida*, *Pyrococcus horikoshii*, *Ralstonia solanacearum*, *Rhizobium leguminosarum*, *Rhodococcus*, *Sinorhizobium meliloti*, *Variovorax paradoxus* and *Zihengliuella alba* [46-52]. Production of the exopolysaccharides (EPS) by PGP microbes plays a vital role in influencing the soil structure. EPS producing microbes stimulate the water binding capacity of soil and help in regulation of supply of nutrients and water to roots. EPS help in irreversible attachment colonization of the microbes to the roots due to network of fibrillar material that permanently connects the microbes to the root surfaces. Bashan, *et al.* [53], demonstrated the role of polysaccharides producing *Azospirillum* in aggregation of the soil. The production of the extracellular biofilms by PGP microbes for binding and making the water molecules in the rhizospheric region available is another strategy for alleviation of the water stress conditions [54]. EPS production has been reported in *Pseudomonas aeruginosa*, *Bacillus subtilis* and *Streptococcus mutans* [55].

Another important consequence of the drought is stimulation of the production of various reactive oxygen species (ROS) including hydrogen peroxide (H_2O_2), singlet oxygen (1O_2), superoxide radical (O_2^-), and the hydroxyl radical ($HO\cdot$) [56] and these reactive oxygen species decreases the normal, metabolic processes of the plants by causing a oxidative damage to the lipids, various proteins ultimately leading to the cell death [57,58]. Plants possess certain enzymatic and non-enzymatic oxidants which are also referred to as the scavenging enzymes which play an efficient and supportive role to overcome the negative effects of the drought [56]. Superoxide dismutase (SOD), catalase (CAT), peroxidase (POX), glutathione reductase (GR), and ascorbate peroxidase (APX) are among the most important enzymatic antioxidants [56-59].

Conclusion and Future Scope

The improvement of the stress tolerance and productivity of the crops is the major goal of agriculture. PGP microbes is an emerging field of science which is proving its potential helping the plants to combat with the abiotic stresses by different mechanisms including production of the phytohormones, solubilization of phosphorus, production of ACC deaminase, production of siderophores. Plant growth-promoting microbe can affect plant growth directly or indirectly. The direct promotion of plant growth by PGP microbes, for the most part, entails providing the plant with a compound that is synthesized by the bacterium or facilitating the uptake of certain nutrients from the environment. The indirect promotion of plant growth occurs when PGP microbes decrease or prevent the deleterious effects of one or more phytopathogenic organisms. Future research in microbes will rely on the development of molecular and biotechnological approaches to increase our knowledge of microbes and to achieve an integrated management of microbial populations of microbial community.

Acknowledgement

The authors duly acknowledge the Department of Biotechnology, Akal College of Agriculture, Eternal University and Vice Chancellor, Eternal University for providing the motivation and research infrastructure.

Bibliography

1. Verma P., *et al.* "Molecular diversity and multifarious plant growth promoting attributes of Bacilli associated with wheat (*Triticum aestivum* L.) rhizosphere from six diverse agro-ecological zones of India". *Journal of Basic Microbiology* 56.1 (2016): 44-58.
2. Yadav AN., *et al.* "Microbiome in Crops: Diversity, distribution and potential role in crops improvements". In: Prasad R, Gill SS, Tuteja N (editors) *Crop Improvement through Microbial Biotechnology* (2018): 305-332.
3. Yadav AN., *et al.* "Actinobacteria from Rhizosphere: Molecular Diversity, Distributions and Potential Biotechnological Applications". In: Singh B, Gupta V, Passari A (editors) *New and Future Developments in Microbial Biotechnology and Bioengineering* (2018): 13-41.
4. Vinocur B., *et al.* "Recent advances in engineering plant tolerance to abiotic stress: achievements and limitations". *Current Opinion in Biotechnology* 16.2 (2005): 123-132.
5. Naveed M., *et al.* "Increased drought stress resilience of maize through endophytic colonization by *Burkholderia phytofirmans* PsJN and *Enterobacter* sp. FD17". *Environmental and Experimental Botany* 97 (2014): 30-39.
6. Lesk C., *et al.* "Influence of extreme weather disasters on global crop production". *Nature* 529.7584 (2016): 84-87.
7. Glick BR., *et al.* "Promotion of plant growth by ACC deaminase-producing soil bacteria". *European Journal of Plant Pathology* 119.3 (2007): 329-339.
8. Yang J., *et al.* "Rhizosphere bacteria help plants tolerate abiotic stress". *Trends in Plant Science* 14.1 (2009): 1-4.
9. Hayat R., *et al.* "Soil beneficial bacteria and their role in plant growth promotion: a review". *Annals of Microbiology* 60.4 (2010): 579-598.

10. Mapelli F, *et al.* "Potential for plant growth promotion of rhizobacteria associated with *Salicornia* growing in Tunisian hypersaline soils". *BioMed Research International* (2013): 248078.
11. Mapelli F, *et al.* "Mineral-microbe interactions: biotechnological potential of bioweathering". *Journal of Biotechnology* 157.4 (2012): 473-481.
12. Yadav AN., *et al.* "Biodiversity and biotechnological applications of halophilic microbes for sustainable agriculture". *Journal of Applied Biology and Biotechnology* 6.1 (2018): 48-55.
13. Nelsen CE., *et al.* "Increased drought tolerance of mycorrhizal onion plants caused by improved phosphorus nutrition". *Planta* 154.5 (1982): 407-413.
14. Raghothama K. "Phosphate acquisition". *Annual Review of Plant Biology* 50.1 (1999): 665-693.
15. Hinsinger P. "Bioavailability of soil inorganic P in the rhizosphere as affected by root-induced chemical changes: a review". *Plant and Soil* 237.2 (2001): 173-195.
16. Yadav AN., *et al.* "Beneficial microbiomes: Biodiversity and potential biotechnological applications for sustainable agriculture and human health". *Journal of Applied Biology and Biotechnology* 5.6 (2017): 45-57.
17. Yadav AN., *et al.* "Plant Growth Promoting Bacteria: Biodiversity and Multifunctional Attributes for Sustainable Agriculture". *Advance in Biotechnology and Microbiology* 5.5 (2017): 1-16.
18. Yadav AN., *et al.* "Current applications and future prospects of eco-friendly microbes". *EU Voice* 3.1 (2017).
19. Son H-J., *et al.* "Solubilization of insoluble inorganic phosphates by a novel salt- and pH-tolerant *Pantoea agglomerans* R-42 isolated from soybean rhizosphere". *Bioresource Technology* 97.2 (2006): 204-210.
20. Wakelin SA., *et al.* "Phosphate solubilization by *Penicillium* spp. closely associated with wheat roots". *Biology and Fertility of Soils* 40.1 (2004): 36-43.
21. Yadav AN., *et al.* "Phosphorus Solubilizing Endophytic Microbes: Potential Application for Sustainable Agriculture". *EU Voice* 2 (2016): 21-22.
22. Yadav AN., *et al.* "Haloarchaea endowed with phosphorus solubilization attribute implicated in phosphorus cycle". *Scientific Reports* 5 (2015): 12293.
23. Yadav AN., *et al.* "Prospecting cold deserts of north western Himalayas for microbial diversity and plant growth promoting attributes". *Journal of Bioscience and Bioengineering* 119.6 (2015): 683-693.
24. Yadav AN., *et al.* "Culturable diversity and functional annotation of psychrotrophic bacteria from cold desert of Leh Ladakh (India)". *World Journal of Microbiology and Biotechnology* 31.1 (2015): 95-108.
25. Mathur N., *et al.* "Mycorrhiza: A Potent Bioinoculant for Sustainable Agriculture". In: Pathak H, Sharma A (editors) *Microbial Technology "The Emerging Era"* Lap Lambert Academic Publishing Ag and Co. Kg, Dudweiler Landstr, Germany (2011): 230-245.
26. Singh RN., *et al.* "First, High quality draft genome sequence of a plant growth promoting and Cold Active Enzymes producing psychrotrophic *Arthrobacter agilis* strain L77". *Standards in Genomic Sciences* 11.1 (2016): 54.
27. Yadav AN. "Bacterial diversity of cold deserts and mining of genes for low temperature tolerance". Ph.D. Thesis, Indian Agricultural Research Institute, New Delhi and Birla Institute of Technology, Ranchi 234 (2015).
28. Yadav AN., *et al.* "Archaea endowed with plant growth promoting attributes". *EC Microbiology* 8.6 (2017): 294-298.
29. Behera B., *et al.* "Diversity, mechanism and biotechnology of phosphate solubilising microorganism in mangrove-a review". *Biocatalysis and Agricultural Biotechnology* 3.2 (2014): 97-110.
30. Goldstein AH "Bioprocessing of rock phosphate ore: essential technical considerations for the development of a successful commercial technology". In: Proceedings of the 4th International Fertilizer Association Technical Conference, IFA, Paris, (2000).
31. Gaba S., *et al.* "Draft genome sequence of *Halolamina pelagica* CDK2 isolated from natural salterns from Rann of Kutch, Gujarat, India". *Genome Announcements* 5.6 (2017): e01593-e01616.
32. Chung H., *et al.* "Isolation and characterization of phosphate solubilizing bacteria from the rhizosphere of crop plants of Korea". *Soil Biology and Biochemistry* 37.10 (2005): 1970-1974.
33. Yadav AN., *et al.* "Extreme cold environments: A suitable niche for selection of novel psychrotrophic microbes for biotechnological applications". *Advances in Biotechnology and Microbiology* 2.2 (2017): 1-4.
34. Saxena AK., *et al.* "Biotechnological applications of microbes isolated from cold environments in agriculture and allied sectors". In: International Conference on "Low Temperature Science and Biotechnological Advances", Society of low temperature biology (2015): 104.
35. Verma P., *et al.* "Growth promotion and yield enhancement of wheat (*Triticum aestivum* L.) by application of potassium solubilizing psychrotolerant bacteria". In: Proceeding of 56th Annual Conference of Association of Microbiologists of India and International Symposium on "Emerging Discoveries in Microbiology" (2015).
36. Verma P., *et al.* "Microbes mediated biofortification of wheat (*Triticum aestivum* L.) for micronutrients by Fe-chelating and Zn-solubilizing bacteria". In: Proceeding of National Conference on Advances in Food Science and Technology (2017).
37. Rana KL., *et al.* "Biotechnological applications of endophytic microbes associated with barley (*Hordeum vulgare* L.) growing in Indian Himalayan regions". In: Proceeding of 86th Annual Session of NASI and Symposium on "Science, Technology and Entrepreneurship for Human Welfare in The Himalayan Region", (2016): 80.
38. Kour D., *et al.* "Biofertilizers: Eco-friendly Technologies and Bioresources for Sustainable Agriculture". In: Proceeding of International Conference on Innovative Research in Engineering Science and Technology, (2017).
39. Srivastava AK., *et al.* "Diversity analysis of *Bacillus* and other predominant genera in extreme environments and its utilization in Agriculture" (2014).
40. Verma P., *et al.* "Alleviation of cold stress in wheat seedlings by *Bacillus amyloliquefaciens* IARI-HHS2-30, an endophytic psychrotolerant K-solubilizing bacterium from NW Indian Himalayas". *National Journal of Life Sciences* 12.2 (2015): 105-110.

41. Solano BR., *et al.* "Physiological and molecular mechanisms of plant growth promoting rhizobacteria (PGPR)". *Plant-bacteria interactions: strategies and techniques to promote plant growth*. Wiley, Weinheim, Germany (2008): 41-52.
42. Whipps JM. "Microbial interactions and biocontrol in the rhizosphere". *Journal of Experimental Botany* 52.1 (2001): 487-511.
43. da Silva V., *et al.* "Rhizobias performance with rhizobacteria growth promoter in plants in the cowpea crop (*Vigna unguiculata* [L.] Walp.)". *Acta Scientiarum-Agronomy* 28.3 (2006): 407-412.
44. Araujo FFd. "Seed inoculation with *Bacillus subtilis*, formulated with oyster meal and growth of corn, soybean and cotton". *Ciência e Agrotecnologia* 32.2 (2008): 456-462.
45. Saikia J., *et al.* "Alleviation of drought stress in pulse crops with ACC deaminase producing rhizobacteria isolated from acidic soil of Northeast India". *Scientific Reports* 8.1 (2018): 3560.
46. Jha B., *et al.* "The roots of the halophyte *Salicornia brachiata* are a source of new halotolerant diazotrophic bacteria with plant growth-promoting potential". *Plant and Soil* 356.1-2 (2012): 265-277.
47. Fujino A., *et al.* "Structural and enzymatic properties of 1-aminocyclopropane-1-carboxylate deaminase homologue from *Pyrococcus horikoshii*". *Journal of Molecular Biology* 341.4 (2004): 999-1013.
48. Gontia I., *et al.* "*Brachybacterium saurashtrense* sp. nov., a halotolerant root-associated bacterium with plant growth-promoting potential". *International Journal of Systematic and Evolutionary Microbiology* 61.12 (2011): 2799-2804.
49. Madhaiyan M., *et al.* "Regulation of ethylene levels in canola (*Brassica campestris*) by 1-aminocyclopropane-1-carboxylate deaminase-containing *Methylobacterium fujisawaense*". *Planta* 224.2 (2006): 268-278.
50. Gontia-Mishra I., *et al.* "Molecular diversity of 1-aminocyclopropane-1-carboxylate (ACC) deaminase producing PGPR from wheat (*Triticum aestivum* L.) rhizosphere". *Plant and Soil* 414.1-2 (2017): 213-227.
51. Yadav AN "Studies of Methylo-trophic Community from the Phyllosphere and Rhizosphere of Tropical Crop Plants". M.Sc. Thesis, Bundelkhand University (2009): 66.
52. Verma P., *et al.* "Appraisal of diversity and functional attributes of thermotolerant wheat associated bacteria from the peninsular zone of India". *Saudi Journal of Biological Sciences* (2016).
53. Bashan Y., *et al.* "Azospirillum-plant relationships: physiological, molecular, agricultural, and environmental advances (1997-2003)". *Canadian Journal of Microbiology* 50.8 (2004): 521-577.
54. Timmusk S., *et al.* "Plant root associated biofilms: perspectives for natural product mining". In: *Bacteria in Agrobiol-ogy: Plant Nutrient Management*. Springer (2011): 285-300
55. Vimala P., *et al.* "Characterization of exopolysaccharide (EPS) produced by *Leuconostoc* sp. V 41". *Asian Journal of Microbiology, Biotechnology and Environmental Sciences* 5.2 (2003): 161-165.
56. Cruz de Carvalho MH. "Drought stress and reactive oxygen species: production, scavenging and signaling". *Plant Signaling and Behavior* 3.3 (2008): 156-165.
57. Farooq M., *et al.* "Plant drought stress: effects, mechanisms and management". In: *Sustainable agriculture*. Springer (2009): 153-188.
58. Hasanuzzaman M., *et al.* "Drought stress responses in plants, oxidative stress, and antioxidant defense". *Climate Change and Plant Abiotic Stress Tolerance* (2013): 209-250.
59. Gill SS., *et al.* "Reactive oxygen species and antioxidant machinery in abiotic stress tolerance in crop plants". *Plant Physiology and Biochemistry* 48.12 (2010): 909-930.

Volume 2 Issue 6 June 2018

© All rights are reserved by Ajar Nath Yadav and Neelam Yadav.