



Study on Chemical Composition, Digestibility and Ruminal Degradation Parameters of Siris Leaves, Flowers and Pods in One-Humped Camel

Tahereh Mohammadabadi^{1*}, Morteza Chaji¹, Zeinab Uosefi¹ and Reza Tahmasbi²

¹Department of Animal Science, Khuzestan Agricultural Sciences and Natural Resources University, Iran

²Department of Animal Science, Shahid Bahonar University of Kerman, Iran

*Corresponding Author: Tahereh Mohammadabadi, Department of Animal Science, Khuzestan Agricultural Sciences and Natural Resources University, Iran.

Received: January 19, 2018; Published: May 28, 2018

Abstract

The aim of this experiment was to determine the chemical composition, digestibility, fermentation and ruminal degradation parameters of siris (*Albizia lebbek*) leaves, flowers and pods by total microorganisms or fungi + protozoa mixture in one-humped camel. Chemical composition of samples was measured and gas production technique and *in situ* method were used for determination of fermentation and rumen degradation parameters, respectively. The result showed the CP values of siris flowers were the greatest whenever NDF, ADF and ether extract of pods were more than leaves and flowers ($p < 0.05$). The gas production potential of siris leaves, flowers and pods by the total microorganisms of camel rumen were 48.1, 22.2 and 51.2 ml, respectively ($p > 0.05$). The lowest gas production potential by fungi + protozoa mixture was recorded for leaves ($p < 0.05$). The difference in gas production rate of the leaves, flowers and pods by total microorganisms, and fungi + protozoa after 96 h incubation was not significant ($p > 0.05$). There was no significant difference among siris leaves, flowers and pods for the partitioning factor, microbial biomass and biomass efficiency by total microorganisms and fungi + protozoa mixture. The rapidly degradable fraction in flowers and pods were different from that in leaves, with siris flowers having the highest rapidly degradable fraction ($p < 0.05$). Potential and effective degradability of siris flowers were higher as compared to leaves and pods ($p < 0.05$). The slowly degradable fraction in pods was lower than in leaves and flowers ($p < 0.05$). Therefore, according to proper rumen degradation and fermentation; siris leaves, flowers and pods especially flower can be used in one-humped camel nutrition to improve rumen degradation processes, effectively.

Keywords: Camel; Gas Production; Chemical Composition; Rumen; Degradation; Siris

Introduction

The camels obtain all their nutritional demands from pasture forages such C4 plants [1,2] that can be found in all tropical grasslands and have lower digestibility than C3 plants [3]. Rumen cellulolytic enzymes activity in camelids is high that have a special ability to use low quality forages higher than the other ruminants [4,5]. It is reported, one of the most abundant organisms of camel rumen are Bacteroides that has been improved animal digestion [6,7]. There is insufficient information available on the microbial ecology of the camels [8].

Siris (*Albizia lebbek*) is a tropical legume in the *Fabaceae* family. It is one of the most widespread and common species of *Albizia* genus in the world and is called "Woman's Tongue", Koko and Lebeck tree. Siris is a native plant of tropical Africa, Asia and northern Australia. It grows up to 5 m in height, and can produce 100 - 120 kg edible dry matter in one year. The fragrant flowers are greenish-yellow to white (2.5 - 7.5 cm in diameter). The tree produces numerous light grey pods (10 - 30 cm long; 2 - 5 cm wide), and each pod contains 4 - 12 pale brown seeds. Most livestock eat leaves and young twigs of this promising fodder tree readily [9]. Carbohydrates are the major components of siris, and potassium and copper are found to be in the highest and lowest among its mineral, respectively [10,11]. Flowers contain no harmful constituents and have relatively high amounts of N and Ca, but low amount of P in compared with other parts of this plant [9,12]. The amino

acids profile indicated that arginine and lysine are present in large amounts in seeds while glutamic acid and aspartic acid are higher in pods [9,11]. The siris leaves are contain 23 % CP and pods contain 19 % CP and 45% NDF [13]. The seeds CP is more than pods [14]. Due to high DM and low moisture of the leaves, they can be used as protein supplements for formulation of livestock rations [14].

Economically, siris is an important industrial and medicinal plant [11]. Leaves and seeds are used for treatment of eye inflammations and the flowers are used for the treatment of spermatorrhea [11]. Siris leaves has a low tannins and phenolic compounds contents. Linoleic acid is the major fatty acid in leaves and pods [11,15]. Total tannin content of siris leaves is around 4 % DM that cannot be having negative effect on healthy of livestock [15]. It is concluded that the tannin by 2 to 4 % DM, protect protein from rumen degradation and increase essential amino acids uptake in the intestine, but tannin content more than 5 %, considerably decrease palatability, feed intake and digestibility [27]. This plant has been used in cow, buffalo, goat and sheep feeding in the different experiments (*in vitro* and *in vivo*).

Therefore, the aim of this experiment was to study chemical composition, fermentation and rumen degradation parameters of siris leaves, flowers and pods by total microorganisms and or fungi+protozoa mixture in one-humped camel.

Materials and Methods

Different parts of 5 siris trees (leaves, flowers and pods) were collected from field of Khuzestan Agricultural Sciences and Natural Resources University, and then milled properly. The sampling was done in plots with dimensions of 100 * 100 meters. Chemical composition of siris leaves, pods and flowers were determined by standard methods of AOAC [16].

Gas production (GP) experiments were done in 3 runs and 4 replicates (syringes) per samples. The rumen fluid was collected and mixed before morning feeding from two fistulated one-humped camels, which was fed for 1 month with a forage based diet (60% straw and 40% alfalfa) and some siris branches for adaptation.

Rumen content was strained through four layers of cheese cloth. The strained and free feed residual of rumen fluid were used as total rumen microbiota. The isolation of ruminal fungi+protozoa were carried out from strained rumen fluid by using antibacterial agents (streptomycin sulfate, potassium penicillin G, and chloramphenicol), which were added at the rate of 0.1 ml per 1 ml to the gas production medium [17].

About 300 ± 10 mg sample (1.0 mm screen, milled) were incubated with 35 ml buffered rumen fluid under continuous CO₂ reflux in 100 ml calibrated glass syringes for 2, 4, 6, 8, 10, 12, 16, 24, 48, 72 and 96h, in a water bath at 39°C [18]. The samples were incubated together with three syringes containing only incubation medium (as blank). After 96 hours of incubation, the content of each syringe was used for determination of ammonia-N (NH₃-N) concentration (Kjeltec 2300 Autoanalyzer, Foss Tecator AB, Hoganas, Sweden). Cumulative gas production data were fitted to the exponential equation $Y=b(1-e^{-ct})$, where b is the gas production (ml) from the fermentable fraction, C is the rate constant of gas production (ml/h), t is the incubation time (h) and Y is the volume of gas produced at time [18].

For determination of the partitioning factor (PF) at the end of each incubation period, the content of syringes was transferred into an Erlenmeyer flask, mixed with 20 ml neutral detergent fiber solution, boiled for 1 hour, filtered, dried (in oven at 60°C for 48 h) and ashed (in furnace, at 550°C for 3h). The partitioning factor, microbial biomass and actual degradable organic matter were calculated by the method of Makkar and Becker.

Dry matter degradability was measured by *in situ* technique using 2 male one-humped camels fitted with rumen fistula (400 ± 12 Kg, BW). Five g (DM basis) of each milled sample (2.0 mm screen) was transferred into a polyester bag (10 × 20 cm, 52 µm pore size) and incubated in the rumen for 2, 4, 6, 8, 16, 24, 48, 72 and 96 hours

(n = 4). At the end of each incubation period, the bags were immediately hand-rinsed under cold tap water until clear and dried in a forced-air oven at 60°C for 48 hours. The bags without incubation (0 h) were washed to estimate the wash-out at initial time. The disappearance of DM and CP were calculated using the equation $P = a + b(1 - e^{-ct})$. Where, P is fraction degraded in the time t, a is soluble fraction, b is potentially degradable fraction, c is degradation rate and t is incubation time. The effective degradability ($k = 0.03, 0.05$ and $0.08/h$) was calculated using the equation $ED = a + (bc/(c+k))$ in which k is the estimated rate of outflow from the rumen [19].

The data were subjected to analysis of variance as a completely randomized design using the General Linear Model (GLM) procedure of the SAS. The Duncan’s multiple range test was used to compare the mean difference at $p < 0.05$.

Results and Discussion

The result of chemical composition of different parts of siris was given in figure 1. The CP value of siris flower was the greatest in compared with pods and leaves whenever NDF, ADF and ether extract of pods were more than leaves and flowers ($p < 0.05$).




Figure 1: Chemical composition of siris leaves, pods and flowers (%).

The potential of gas production by total rumen microorganisms were 48.07, 22.71 and 51.16 ml/300 mg for leaves, flowers and pods, respectively ($p > 0.05$). However, the corresponding values by fungi+protozoa mixture were 11.54, 29.65 and 18.91 ml, respectively (Table 1). No significant difference was observed in potential and gas production rate during 96 h incubation period for leaves, pods and flowers of siris in mixed fungi + protozoa ($p > 0.05$) (Table 2). There was no significant difference in PF amount and microbial biomass in the presence of two microbial groups ($p > 0.05$) (Table 1 and 2).

	Partitioning factor (mg mL ⁻¹)	Microbial biomass (mg)	Microbial biomass efficiency (%)	Actual organic matter disappearance (mg)	Cell wall degradation (%)	b (mL)	c(mL/h)
Leaf	3.50	24.30	36.83	46.20	0.76	48.07	0.03
Pod	3.33	115.35	33.58	39.45	0.20	22.71	0.05
Flower	2.54	88.95	17.39	11.95	0.53	51.16	0.04
SEM	0.23	47.42	11.32	18.41	0.71	15.70	0.01
P-value	0.10	0.10	0.10	0.10	0.10	0.10	0.50

Table 1: Gas production parameters of siris leaves, pods and flowers by total rumen microorganisms b: gas production from the fermentable fraction, c: rate constant of gas production. SEM: Standard error of the mean.

	Partitioning factor (mg mL ⁻¹)	Microbial biomass (mg)	Microbial biomass efficiency (%)	Actual organic matter disappearance (mg)	Cell wall degradation (%)	b(mL)	c (mL/h)
Leaf	2.90	25.95	24.37	106.25	0.33	11.54	0.07
Pod	2.77	18.10	20.42	87.40	0.36	29.65	0.02
Flower	2.43	11.95	12.88	90.55	0.40	18.91	0.06
SEM	0.51	8.41	7.31	11.86	0.65	9.66	0.05
P-value	0.10	0.10	0.10	0.10	0.10	0.10	0.90

Table 2: Gas production parameters of siris leaves, pods and flowers by fungi+protozoa mixture in one - humped camel.

SEM: Standard error of the mean.

The effects of siris leaves, pods and flowers on ammonia nitrogen in gas environment (Table 3) were significant. Siris flowers in presence of the two microbial groups had maximum concentrations of ammonia nitrogen ($p < 0.05$).

The difference between rapidly degradable fraction of DM in pods and flowers was significant (Table 4). The potential and effective degradability of siris flowers increased significantly as compared to leaves and pods ($p < 0.05$). The slowly degradable fraction of pods decreased more significantly than leaves and flowers. The rapid degradable fraction of flowers and pods as compared to siris leaves was considerably different. Furthermore, potential and effective degradability of siris flower in comparison to leaf and pod were increased significantly (Table 4).

	Total microorganisms		Fungi + protozoa	
	Ammonia Nitrogen (mg/ 100 mL)	pH	Ammonia Nitrogen (mg/ 100 mL)	pH
Leaf	19.77 ^b	7.18	14.60 ^a	7.25
Pod	14.85 ^c	7.22	10.38 ^b	7.27
Flower	22.72 ^a	6.75	15.65 ^a	6.65
SEM	0.60	0.14	0.30	0.26
P-value	0.006	0.10	0.002	0.30

Table 3: Fermentation parameters of siris leaves, pods and flowers by rumen microorganism in one-humped camel

SEM: Standard error of the mean, ^{a,b}: Means with common letter (s) within each column do not differ significantly ($p > 0.05$).

	Rapidly degradable	Slowly degradable	Constant rate of degradation	Potential degradability	Effective degradability
Leaf	0.31 ^b	0.39 ^a	0.08 ^a	0.70 ^b	0.60 ^b
Pod	0.35 ^{ab}	0.20 ^b	0.06 ^a	0.56 ^c	0.49 ^c
Flower	0.43 ^a	0.42 ^a	0.03 ^b	0.87 ^a	0.67 ^a
SEM	0.02	0.01	0.007	0.008	0.003
P-value	0.03	0.0002	0.01	0.0001	0.0001

Table 4: Parameters of in situ dry matter rumen degradation of siris leaves, pods and flowers in one-humped camel.

SEM: Standard error of the mean, ^{a,b}: Means with common letter (s) within each column do not differ significantly ($p > 0.05$).

Siris flowers have higher protein and ether extract, and lower NDF and ADF in compared to leaves and pods. Flowers are used as a valuable protein supplement due to the high digestible nitrogen. The value of pods NDF and ADF were 56.1% and 42.1%, respectively [20]. The researchers [21] reported NDF, ADF, and ether extract content of leaves were 46.9, 33.7 and 5.4 %, respectively. But the others [22] evaluated the content of NDF, ADF, and ether extract of leaves were 48.44, 36.87 and 3.62 %, respectively.

Researches indicated the crude protein content of the siris leaves of the region of Townsville in Australia was 17.5% [23]. Also, it is reported that siris leaves can be used as a protein supplement instead of cotton seed meal to feed the goats [24]. In another experiment, the protein content of *Albizia julibrissin* and *Albizia procera* leaves were reported as 18.61% and 17-24%, respectively [25,26]. The reason for the difference in chemical composition depends on the differences in the sampling season, plant growth stage, species type, climatic differences and conditions in the areas [22].

Based on the result, gas production potential and rate of different parts of siris by total rumen microorganisms were not significant. But in contrast with this results, researchers reported that presence of tannin in siris leaves reduced gas production, because of tannins affinity to make a complex with nutrients, therefore, it keeps nutrients away from microorganisms and prevents their fermentation [27].

Tannin content in the siris leaves and seed was reported about 4 and 5.3%, respectively [15]. Tannins can also reduce microorganism adhesion to nutrients, inhibit microorganism growth and microbial enzymes activity which have negative effects on fermentation, nutrients digestibility and methane production [28].

The flowers are free of antinutrient and toxic compounds [11]. It is concluded, lack of anti-nutritional complex and higher quantities of proteins and carbohydrates in flowers of siris, makes it as suitable substrate for rumen microbial growth [9]. Also, they have higher gas production than leaves and pods because of lower NDF

and ADF [13]. It is reported, decreasing the content of lignin in the plant cell wall enhanced the fermentation and gas production [30]. Similar to the current findings, it was reported that when the siris flower was fed by 15 - 16%, the dry matter digestibility of the forage with the flower increased [29].

The mount of PF and microbial biomass in the two microbial groups were not different. It was reported that the tannin and saponin of leaves and pods of siris increases the PF [11].

The amount of saponin in siris branches (pods) was 669.4 mg/kg, that was higher than other anti-nutritional agents. It is suggested that the pods be separated from the seeds in order to increase the nutritional value. Also, tannin content in the siris leaves and pods mixture was about 4 % [11].

Generally, the feedstuff that contains tannin, have higher PF which might be related to tannins being dissolved during fermentation and decrement of dry matter. However, the latter had no effect on gas production or microbial protein synthesis. Tannins become complexed with proteins, the presence of which in the undigested residues results in under-estimation of the actual digested organic matter. Therefore, an error in PF quantity will occur [28]. The researchers reported that presence of resources which contains tannin, increased PF [31]. They found that tannin has a positive effect on protein nutrition in animal feeding. Digested nutrients are more effective in microbial protein synthesis as compared to short chain fatty acid synthesis. Siris belongs to *Fabaceae*, so improving those parameters can be related to legume ability to provide nitrogen requirement, energy and vitamins for microbial growth [32].

Saponin can increase microbial protein synthesis in vitro due to its negative effect on protozoa which prevents bacterial engulfing by protozoa and increases microbial nitrogen flow to the duodenum [18]. Moreover, it was reported that PF in some legume plants like *Leucaena leucocephala* and *acacia* (*Fabaceae* family) were 4.12 and 3.78, respectively [30].

Siris flowers in presence of two microbial groups had maximum concentrations of ammonia nitrogen which might be related to their higher protein value (i.e. 22.75%). Reduction of ammonia nitrogen concentration in siris pods might be related to the presence of saponin, and its negative effect on protozoa population. Therefore, by reduction of protozoa, bacteria lysis decreases and less ammonia will be produced, and ammonia concentration reduces indirectly [33]. Although, ammonia will be produced in the rumen through bacterial degradation by protozoa [33]. Protozoa has proteolytic and deamination effect which cause ammonia production in rumen [34]. By reducing protozoa in rumen, nitrogen flow between bacteria and protozoa reduces and nitrogen flow from the rumen increases [33]. Protozoa population reduction in rumen and decrement of ammonia nitrogen concentration might be due to negative effect of saponin on protozoa population [35].

The researchers reported that resources which contain tannins have reducing effect on ammonia nitrogen concentration in the rumen [28]. The effects of tannins on protein metabolism in the rumen might be related to tannins ability to attach to protein in order to reduce microbial enzyme activity, proteolytic bacteria growth and consequently protein degradation and ammonia nitrogen production in the rumen decreased. Another reason for reducing pH in diet which contains leaves, might be protozoa population

reduction. Rapid digestion and storage of starch by protozoa cause stable rumen and constant pH [15].

Higher value of rapidly degradable fraction of siris flowers as compared to pods and leaves might be due to their higher protein content, soluble carbohydrate and lower ADF and NDF contents [36]. It seems that increasing crude protein content of siris had positive effect on DM degradation. By increasing protein content in plant, DM potential of degradability will be increased. Generally, degradability is affected by cell wall [37]. Significant reduction of slowly degradable fraction of pods as compared to leaves and flowers of siris might be related to higher values of NDF and lignin in pod [28].

On the base of studies [38] digestibility of dry matter, organic matter and crude fiber and nitrogen retention of siris leaves were 89.87%, 93.9%, 61.44% and 91.16% respectively. Also, it is concluded that siris silage could be used as source of protein for sheep fed with low-quality rations [38].

Presence of anti-nutritional factors such as saponins and oxalate in siris pod might be another reason for lower slowly degradable fraction, as these compounds reduce fiber digestibility in rumen. Saponins suppress fiber digestion in rumen, which might be related to reducing effect of fibrolytic enzyme activity [39]. Potential and effective degradability of siris flower in comparison to leaf and pods were significantly increased. It might be due to the presence of anti-nutritional factors like saponin and oxalate in siris leaves and pods.

The researchers reported the rapidly degradable fraction, slowly degradable fraction, degradation rate constant and potential of degradability of siris leaves were 42.2, 37.3, 0.1 and 79.5%, respectively [40]. It is seen that the rapidly degradable fraction and potential of degradability of the present research is lower, but slowly degradable fraction are same [11].

Conclusion

It can be concluded that siris leaves, flowers and pods have good rumen degradability in one-humped camel. Feeding camel with branches of this plant especially flower increase fiber digestion through improvement of rumen fermentation processes.




Figure 2: Gas production kinetics of siris leaves, pods and flowers by total rumen microorganisms in one-humped camel.




Figure 3: Gas production kinetics of siris leaves, pods and flowers by fungi + protozoa mixture in one-humped camel.

Acknowledgements

The authors gratefully acknowledge the khuzestan Agricultural Sciences and Natural Resources University for their laboratory and financial support.

Conflict of Interest

There is no conflict on any issues relating to the research between the authors.

Bibliography

- Samsudin AA, et al. "Molecular diversity of the foregut bacteria community in the dromedary camel (*Camelus dromedarius*)". *Environmental Microbiology* 13.11 (2011): 3024-3035.
- Ahmed MH, et al. "Effect of energy source supplementation of the utilization of some desert forage by growing Lambs". *Journal Advance Agriculture Resource Resources* 6 (2001): 255-277.
- Haddi ML, et al. "Seasonal changes in chemical composition and in vitro gas production of six plants from Eastern Algerian arid regions". *Livestock Research Rural Development* 21.4 (2009).
- Robinson TF, et al. "Digestibility and nitrogen retention in llamas and goats fed alfalfa, C3 grass, and C4 grass hays". *Small Ruminant Research* 64.1-2 (2006): 162-168.
- Jouany JP. "Defaunation of the rumen. In: *Rumen Microbial Metabolism and Ruminant Digestion*" (1991).
- Bhatta R, et al. "Effect of supplementation containing polyethylene glycol (PEG)-6000 on intake, rumen fermentation pattern and growth in kids fed foliage of *Prosopis cineraria*". *Small Ruminant Research* 52.1-2 (2004): 45-52.
- Miron J, et al. "Invited review: adhesion mechanisms of rumen cellulolytic bacteria". *Journal Dairy Science* 84.6 (2001): 1294-1309.
- Ghali MB, et al. "Identification and characterization of the predominant lactic acid-producing and lactic acid-utilizing bacteria in the foregut of the feral camel (*Camelus dromedarius*) in Australia". *Animal Production Science* 51.7 (2011): 597-604.
- Lowery JB. "Agronomy and forage quality of *Albizia lebbek* in the semi-arid tropics". *Tropical Grasslands* 23.2 (1989): 84-91.
- Mozafarian V. "Dicotyledon Classification". Amir Kabir Publications 4th edition (2005): 620.
- Zia UM, et al. "Compositional studies and antioxidant potential of *Albizia lebbek* (L.) Benth. Pods and seeds". *Turkish Journal Biology* 37(2013): 25-39.
- Gupta BS. "Nutritive value of siris (*Albizia lebbek*) tree leaves". *Indian Journal of Nutrient Diet* 17 (1980): 187-191.
- Kennedy Peter M, et al. "Utilization of tropical dry season grass by ruminants is increased by feeding fallen leaf of siris (*Albizia lebbek*)". *Animal Feed Science and Technology* 96.3 (2002): 175-192.
- Hassan K, et al. "Nutritional Evaluation of *Albizia lebbek* (L.) Pods as Source of Feeds for Livestock". *American Journal of Food Technology* 2.5 (2007): 435-439.
- El-Hawary K, et al. "A phytochemical profile of *Albizia lebbek* L. Benth. cultivated in Egypt". *Asian Journal of Biochemistry* 6.2 (2011): 122-141.
- Association of Official Analytical Chemists. Official Method of Analysis 15th edition AOAC Arlington (2002).
- Zhang Y, et al. "Fermentation of plant cell walls by ruminal bacteria, protozoa and fungi and their interaction with fibre particle size". *Archive Animal Nutrition* 61.2 (2010): 114 - 125.
- Menke KH, et al. "Estimation of the energetic feed value obtained from chemical analysis and gas production using rumen fluid". *Animal Research Development* 28 (1988): 7-55.
- Orskov ER, et al. "The estimation of protein degradability in the rumen from incubation measurements weighed according to rate of passage". *Journal of Agricultural Science* 92.2 (1979): 499-503.
- Nyambati EM. "The value of *Acacia brevispica* and *Leucaena leucocephala* seedpods as dry season supplements for calves in dry areas of Kenya". *African Journal of Agricultural Research* 1.4 (2006): 118-124.
- Balogun RO, et al. "Digestibility of some tropical browse species varying in tannin content". *Animal Feed Science and Technology* 76.1-2 (1998): 77-88.
- Balgees A. "Effects of *Albizia lebbek* or wheat bran supplementation on intake, digestibility and rumen fermentation of ammoniated bagasse". *Journal of Applied Science Research* 5.8 (2009): 1002-1006.
- Dwatmadji Teleni E, et al. "Nutritive value of *Albizia lebbek* supplements for growing sheep". *Australian Journal of experimental Agriculture* 32 (1992): 273-278.
- Ndemanisho EE, et al. "The potential of *Albizia lebbek* as a supplementary feed for goats in Tanzania". *Agroforestry System* 67.1 (2006): 85-91.
- Alam MR. "Effect of tannins in *Acacia nilotica*, *Albizia procera* and *Sesbania acculeata* foliage determined in vitro, in sacco, and in vivo". *Asian Australian Journal of Animal Science* 20.2 (2007): 220-228.
- Bouazza L, et al. "Nutritive evaluation of foliage from fodder trees and shrubs characteristic of Algerian arid and semi-arid areas". *Journal of Animal and Feed Science* 21.3 (2012): 521-536.

27. McSweeney CS., et al. "Microbial interactions with tannins: nutritional consequences for ruminants". *Animal Feed Science and Technology* 91.1-2 (2001): 83-93.
28. Hassan Sallam SMA., et al. "Ruminal fermentation and tannins bioactivity of some browses using a semi-automated gas production technique". *Tropical and Subtropical Agroecosystem* 12.1 (2010): 1-10.
29. Sommart K., et al. "Fermentation characteristics and microbial protein synthesis in an in vitro system using cassava, rice straw and dried ruzi grass as substrates". *Animal Sciences* 13.8 (2000): 1084-1093.
30. Soltan YA. "Comparative in vitro evaluation of forage legumes (prosopis, acacia, atriplex, and leucaena) on ruminal fermentation and methanogenesis". *Journal of Animal Feed Science* 21.4 (2012): 759 -772.
31. Angaji L., et al. "Deactivation of tannins in raisin stalk by polyethylene glycol-600: Effect on degradation and gas production in vitro". *African Journal of Biotechnology* 10.21 (2011): 4478-4483.
32. Hassoun P., et al. "Feeding dairy heifers untreated or urea treated fibrous sugarcane residues effect on dry matter intake, growth and metabolic parameters". *Animal Feed Science and Technology* 100.1 (2002): 31-41.
33. Wina E., et al. "Saponins containing methanol extract of *Sapindus rarak* affect microbial fermentation, microbial activity and microbial community structure in vitro". *Animal Feed Science and Technology* 121.1 (2005): 159-174.
34. Williams AG., et al. "The Rumen Protozoa". Springer-Verlag, Inc. New York, NY, USA (1991): 441.
35. Wallace RJ., et al. "Natural products as manipulators of rumen fermentation". *Asian-Australians Journal of Animal Sciences* 15.10 (2002): 1458 -1468.
36. Danesh Mesgaran M. "The new in vitro methods in animal researches". Ferdowsi University Press, Mashhad (2009): 191.
37. Arzani H., et al. "Phenological effects on forage quality of five grass species". *Journal of Range Management* 57.6 (2004): 624 -630.
38. Abdulrazak SA., et al. "Nutritive evaluation of *Prosopis juliflora* fruits and leaves from Kenya: Chemical composition and in vitro gas production". In *Proceeding of British Society of Animal Science*, Scarborough 22 (1999): 146.
39. Liu CD., et al. "Alfalfa saponins affect site and extent of nutrient digestion in ruminants". *Journal of Nutrition* 117.5 (1987): 919-927.
40. Larbi A. "Feed value of multipurpose fodder trees and shrubs in West Africa: edible forage production and nutritive value of *Millettia thonningii* and *Albizia lebbek*". *Agroforestry System* 33.1 (1996): 41-50.

Volume 2 Issue 6 June 2018

© All rights are reserved by Tahereh Mohammadabadi, et al.