



Assessment of Genetic Fidelity of Micro Propagated Plants of *Aerva lanata* (L.) Juss. ex Schult. using DNA Markers

Alok Ranjan Sahu^{1,2*} and Jogeswar Panigrahi^{1,3}

¹Plant Biotechnology Laboratory, School of Life Sciences, Sambalpur University, Sambalpur, Odisha, India

²Saraswati +3 Science College, Near Petrol Pump, Kesinga Road, Bhawanipatna, Odisha, India

³Department of Biotechnology, Central University of Rajasthan, Kishangarh, Rajasthan, India

*Corresponding Author: Alok Ranjan Sahu, Principal, Saraswati +3 Science College, Near Petrol Pump, Kesinga Road, Bhawanipatna, Odisha, India.

Received: August 30, 2018; Published: October 26, 2018

Abstract

Aerva lanata (L.) Juss. ex Schult. is a valued medicinal herb. The demand of *A. lanata* is gradually increasing, however the supply is inadequate. Therefore, the need for *in vitro* propagation of this plant is crucial. To check the genetic uniformity and true to true type nature of micro-propagated plants were determined by using DNA markers. Among DNA markers RAPD (random amplified polymorphic DNA) were screened to test the genetic stability of *A. lanata* plants regenerated from nodal segments through direct organogenesis. During the study a total of ten RAPD primers were screened, which produced a total of 32 clear, distinct and monomorphic bands. The micro-propagation protocol developed by our group for rapid *in vitro* multiplication is suitable for clonal propagation and RAPD marker analysis for genetic fidelity assessment of *A. lanata*.

Keywords: Genetic Fidelity; *Aerva lanata*; DNA Markers

Introduction

Aerva lanata (L.) Juss. ex. Schult. is an erect dioecious herb, about 80cm in height having cylindrical, branched tap root belonging to Amaranthaceae family. Shoots of this herb are covered with smooth hairs; leaves are simple and alternate, along with obtuse or acute apex. Further the inflorescence consists of axillary spikes and flowers are bisexual, small sessile, greenish white with spikes [37]. This medicinal herb is commonly known as 'Chhaya' in Hindi, 'Bhadram' in Sanskrit, 'Pulai' in Telugu and 'Lopang Ark' in Odia. It has been reported that this plant is used as anti-inflammatory, diuretic, antimicrobial, anticancer, nephroprotective, antidiabetic, antihelminthic, antifertility activities [1-10]. *A. lanata* is loaded with broad range of phyto-constituents. Phytochemical screening showed the presence of various classes of phytochemicals like alkaloids, amino acids and proteins, carbohydrates, saponins, steroids, flavinoids, and tannins [15,16]. The roots possess good amount of aervin, aervoside and methylaervin [17-19], which possibly contribute to its diverse uses in folklore medicines. It has been reported that the whole plant and plant parts including

roots and leaves has been used in several herbal preparations to cure various ailments in various pharmaceuticals like Bhadraveradi Kashyam, Dasmolam Kashayam, Nishakathakadi Kashayam (www.asoka.co.in), Mahasiddhartha oil (www.aparmita.lv). As the raw plant extracts were rich in antioxidants, hence used as tonic by rural poor during pregnancy [11]. At present this important herb is fulfill its demands only from the natural sources. However, wide use of this herb may cause a possible threat for its survivals. Hence *in vitro* culture techniques like organogenesis and somatic embryogenesis may apply for their propagation and conservations of this therapeutic plant [20-21].

Plant tissue culture technique has long been recognized as an efficient tool for rapid clonal multiplication of true to the type. However, the possibility of developing somaclonal variations still exists [23,24] due to several factors such as genotypes used, composition of growth regulators, expression of chromosomal mosaicism, spontaneous mutation, etc. on the other hand, the

occurrence of stable somaclonal variations of specific type might be advantageous for the crop improvement programme [25,26]. Monitoring the degree of genetic fidelity among micropropagated plants using molecular markers is desirable to reduce the chances for inclusion of variable genotypes and also to assess the technology that is being developed to be marketed. Various molecular markers had been used to assess the genetic fidelity of *in vitro* raised plants by various authors. However, Random Amplified Polymorphic DNA (RAPD) is the cheapest yet reliable and could be a powerful tool for the detection of genetic variability in plants [27-33]. Keeping that entire thing in mind the present objective of this report is assessment of genetic fidelity of micropropagated plants of *A. lanata* using DNA markers.

Materials and Methods

The micro-propagation of *A. lanata* protocol was standardized by Sahu, *et al.* [35], Sahu [36] at Plant Biotechnology Lab, School of Life Sciences, Sambalpur University. For the present study *in vitro* cultures from nodal segments were established on the MS [22] medium fortified with equal conc. BA and kinetin (3.0 mg/l) was found to be the best for direct multiple shoot bud induction, while addition of NAA to the medium with optimum concentration of BA and kinetin fortified MS medium was not found suitable for direct organogenesis. Rather this addition of NAA promotes elongation of shoots and callus mediated organogenesis. For successful rooting of well-developed shoots half strength of MS medium fortified with IBA (0.5mg/L) was used. The survival rate of regenerated plantlets of *A. lanata* was $72 \pm 4\%$ [34,35]. The *in vitro* regenerated plants were morphologically alike with the mother plant. For more evidence RAPD banding pattern was compared between mother plant and nine randomly selected *in vitro* regenerated plants by using RAPD primers.

Isolation and purification of genomic DNA

For genomic DNA isolation, about 0.8g of fresh leaves were ground to fine powder in liquid nitrogen and DNA was isolated following the CTAB method [33] with modifications by adding activated charcoal (0.2% v/v) and PVP (3.5% v/v) to the extraction buffer. DNA concentration and purity were measured by using UV-Vis spectrophotometer (UV-1, Thermo, USA) with T10E1 buffer (pH 8.0) as blank. For further confirmation the quantification of DNA was accomplished by analyzing the purified DNA on 0.8%

agarose gel along with diluted uncut lambda DNA as standard. All the DNA samples were equilibrated to concentration of 25ng/ μ l using T10E1 buffer.

RAPD marker analysis

Ten RAPD primers were used for polymorphism detection. List of primers and their information were presented in Table 1. For RAPD analysis PCR amplification of 25 ng of genomic DNA was carried out using decamer oligonucleotide primers (Operon Tech., Alameda, CA, USA). Each amplification reaction of 25 μ l mixture contained the 25 ng template DNA, 2.5 μ l of 10X assay buffer (100 mM Tris.Cl, pH 8.3; 0.5M KCl; 0.1% gelatin), 2.0mM MgCl₂, 200 μ M each of dNTPs, 20 ng primer, 1.0 units Taq DNA polymerase (Bangalore Genei Pvt. Ltd., Bangalore, India). The amplification was carried out in a thermal cycler (Gene Amp-9700, Applied Biosystem, USA) programmed as initial denaturation at 94°C for 5 minutes, cycle denaturation at 94°C for 2 minutes, cycle annealing at 37°C for 1 minute, cycle elongation at 72°C for 2 min for 45 cycles followed by final elongation at 72°C for 5 minutes.

Amplified products were separated on 1.4% agarose gel containing ethidium bromide solution (@ 0.5 μ g/ml of gel solutions) in submarine electrophoresis apparatus (Tarson-Midi, India) using TAE (40mM Tris acetate; 2mM EDTA) buffer at constant 50 V for about 1-2 hours. A gel loading buffer (20% Sucrose; 0.1 M EDTA, 1.0% SDS; 0.25% Bromophenol blue; 0.25% Xylene cyanol) was used as tracking dye. Amplified products were visualized by Gel documentation system (Geldoc-XR, Bio-Rad, USA) and photographed. The sizes of the amplified products were determined using 250bp ladder (Bangalore Genei Pvt. Ltd., Bangalore, India) as standard and Quantity One software.

Results and Discussion

Genetic fidelity testing of the micropropagated plantlets needs to be authenticated for commercial scale application of the developed micropropagation protocol. PCR-based molecular markers have emerged as simple, fast, reliable and labor-effective tools for testing the genetic fidelity of *in vitro* raised plantlets. Ten RAPD primers of OPA series (OPA01-OPA10) generated 32 clear amplicons in total (Figure 1). The sizes of the amplified bands were ranging from 400 bp to 2,250 bp. A maximum of six bands was generated from OPA-05 while a minimum of one band was

generated from OPA-07 (Table 1). Monomorphic banding pattern was observed for all the amplified band classes across all the micro propagated plants, along with the donor mother plant with all the tested primers. No polymorphism was observed revealing the genetic integrity of *in vitro* regenerated plants. This method often yields large number of markers; it is technically simple to perform and does not need either large amounts of DNA or previous information of DNA sequences, which are often unknown in many plants. Moreover, RAPD analysis is comparatively easy, cost effective and requires comparatively minimum time and also involves no radioactive compound to be handled. Employing RAPD technique, various authors have reported the absence of genetic variation in many species such as *Zingiber officinales* [27], *Chlorophytum borivilianum* [30], *Picea marina* [31], and *Festuca pratensis* [32], using RAPD analysis. As observed, our result clearly indicates the genetic integrity and true-to-type nature of the *in vitro* regenerated plants in germplasm conservation program.

Primer	Primer sequence (5' - 3')	No. of total bands amplified	Size range (bp)
OPA - 01	5'- CAGGCCCTTC-3'	2	855 -1065
OPA - 02	5'- TGCCGAGCTG-3'	3	400 -2250
OPA - 03	5'- AGTCAGCCAC-3'	2	980 -1275
OPA - 04	5'- AATCGGGCTG-3'	5	655 -1945
OPA - 05	5'- AGGGGTCTTG-3'	6	510 -1645
OPA - 06	5'- GGTCCCTGAC-3'	5	645 - 1700
OPA - 07	5'- GAAACGGGTG-3'	3	525 -1755
OPA - 08	5'- GTGACGTAGG-3'	1	710
OPA - 09	5'- GGGTAACGCC-3'	3	485 - 1125
OPA - 10	5'- GTGATCGCAG-3'	2	1125 -1780

Table 1: List of primers, their sequences, number and size of the amplified fragments generated by random amplified polymorphic DNA (RAPD) primers.




Figure 1: Protuberances from the cut end node explant at the end of 2nd week (a), Multiple shoot initials on MS with 3.0 mg/l BA and 3.0mg/l Kn after 8th week (b), Elongated shoot showing origin of roots on MS medium with 0.5 mg/l IBA (c), Regenerated plantlets in plastic cups containing sand and soil mixture during acclimatization (d), and RAPD profiles, M: Mol. weight Ladder, D: donor plant, and 1-9: randomly selected micro propagated plants.

Conclusion

Aerva lanata (L.) Juss. ex Schult. is a valued medicinal herb. The demand of *A. lanata* is all the time on the increase, however the supply is inadequate. Therefore, the need for in vitro propagation of this plant is crucial. To check the genetic uniformity and true to true type nature of micro-propagated plants were determined by using molecular markers. Among DNA markers RAPD (random amplified polymorphic DNA) were screened to test the genetic stability of *A. lanata* plants regenerated from nodal segments through direct organogenesis. During the study a total of ten RAPD primers were screened, which produced a total of 32 clear, distinct and monomorphic bands. The micro-propagation protocol developed by our group for rapid *in vitro* multiplication is suitable for clonal propagation and RAPD marker analysis for genetic fidelity assessment of *A. lanata*.

Acknowledgements

The authors are grateful to the Vice Chancellor, Sambalpur University for financial support and the Head, School of Life Sciences, Sambalpur University for providing necessary facilities. The author (ARS) especially thankful to the Chairman, the Director and the Vikash management groups for managing the Saraswati +3 Science College, Bhawanipatna and for moral supports, and encouragement.

Bibliography

1. Vertichelvan T, *et al.* "Diuretic and anti-inflammatory activities of *Aerva lanata* in rats". *Indian Journal of Pharmaceutical Science* 62 (2000): 300-302.
2. Chowdhury D., *et al.* "Antimicrobial activity and cytotoxicity of *Aerva lanata*". *Fitoterapia* 73.1 (2002): 92-94.
3. Nevin KG and Vijayammal PL. "Effect of *Aerva lanata* on solid tumor induced by DLA cells in mice". *Fitoterapia* 74.6 (2003): 578-582.
4. Shirwaikar A., *et al.* "Effect of *Aerva lanata* on cisplatin and gentamicin models of acute renal failure". *Journal of Ethnopharmacology* 90.1 (2004): 81-86.
5. Venkatesh S., *et al.* "Antinociceptive effect of *Aerva lanata* ethanolic extract in mice: A possible mechanism". *Asian Journal of Pharmacodynamics and Pharmacokinetics*, 9.1 (2009): 58-62.
6. Vetrichelvan T and Jegadeesan M. "Anti-diabetic activity of alcoholic extract of *Aerva lanata* (L.) Juss. Ex Schultes in rats". *Journal of Ethnopharmacology* 80 (2002): 103-107.
7. Nevin KG and Vijayammal PL. "Effect of *Aerva lanata* (L.) Juss against hepatotoxicity of carbon tetrachloride in rats". *Environmental Toxicology and Pharmacology* 20 (2005): 471-477.
8. Appia KG., *et al.* "Hypoglycemic and antihyperlipidaemic effect of ethanolic extract of aerial parts of *Aerva lanata* Linn. in normal and alloxan induced diabetic rats". *International Journal of Pharmaceutical Sciences and Drug Research* 1.3 (2009): 191-194.
9. Anantha A., *et al.* "In vitro antihelmentic activity of aqueous and alcoholic extracts of *Aerva lanata* seeds and leaves". *Journal of Pharmaceutical Sciences and Research* 2.5 (2010): 317-321.
10. Savadi RV and Alagawadi KR. "Antifertility activity of ethanolic extracts of *Plumbago indica* and *Aerva lanata* on albino rats". *International Journal of Green Pharmacy* (2010): 230-233.
11. Mishra L. "Anubhut yogamala and sahaja chikishya" Raj Edition, India (1985).
12. T Pulliah and CK Naidu C. "Antidiabetic plants in India and herbal based antidiabetic research". *Regency Publications* (2003).
13. Pulliah T and Naidu CK. "Antidiabetic plants in India and herbal based antidiabetic research". New Delhi:Regency Publications (2003): 68-69.
14. Muthu C., *et al.* "Medicinal plants used by traditional healers in Kancheepuram District of Tamil Nadu, India". *Journal of Ethnobiology and Ethnomedicine* 2 (2006): 43.
15. Rajadurai M., *et al.* "Ethnomedicinal plants used by the traditional healers of Pachamalai hills, Tamilnadu, India". *Ethno-Medicine* 3.1 (2009): 39-41.
16. Varghese KJ., *et al.* "Dasapushpam: The traditional uses and therapeutic potential of ten sacred plants of Kerala state in India". *International Journal of Pharmaceutical Science and Research* 1 (2010): 50-59.
17. Rajesh R., *et al.* "Antihyperglycemic and antihyperlipidemic activity of aerial parts of *Aerva lanata* Linn Juss in streptozotocin induced diabetic rats". *Asian Pacific Journal of Tropical Biomedicine* 2 (2012): 924-929.

18. Chandra S and Shastri MS. "Chemical constituents of *Aerva lanata*". *Fitoterapia* 61 (1990): 188-190.
19. Afaq SH., et al. "Bisheri Booti (*Aerva lanata*) some lesser known uses and pharmacogonosy". *Ethnobotany* 5 (1991): 37.
20. Zapesochnaya G., et al. "Canthin-6-one and beta-carboline alkaloids from *Aerva lanata*". *Planta Medica* 58 (1992): 192-196.
21. Yuldashev AA., et al. "Components of *Aerva lanata*". *Chemical and Natural Compound* 38 (2002): 293-294.
22. Vasil IK. "Progress in the regeneration and genetic manipulation in crops". *Biotechnology* 6 (1988): 397-402.
23. Murashige T and Skoog F. "A Revised medium for rapid growth and bioassays with tobacco tissue culture". *Physiologia Plantarum* 15 (1962): 473.
24. Rani V and Raina SN. "Genetic fidelity of organized meristem derived micro propagated plants: a critical reappraisal". *In vitro Cellular and Developmental Biology Plant* 36 (2000): 319-330.
25. Bindiya K and Kanwar K. "Random amplified polymorphic DNA (RAPDs) markers for genetic analysis in micro propagated plants of *Robinia pseudoacacia* L". *Euphytica* 135 (2003): 41-47.
26. Karp A. "Somaclonal variation as a tool for crop improvement". *Euphytica* 85 (1995): 295-302.
27. Jain SM., et al. "Somaclonal variation and induced mutations in crop improvement". *Current Plant Science and Biotechnology in Agriculture* 32 (1998).
28. Rout GR., et al. "Determination of genetic stability of micro propagated plants of ginger using random amplified polymorphic DNA (RAPD) markers". *Botanical Bulletin-Academia Sinica Taipei* 39 (1998) 23-27.
29. Rahman MH and Rajora OP. "Microsatellite DNA somaclonal variation in micropropagated trembling aspen (*Populus tremuloides*)". *Plant Cell Report* 20 (2001): 531-536.
30. Phillips RL., et al. "Genetic instability of plant tissue cultures: breakdown of normal controls". *Proceeding of National Academic of Science USA* 91 (1994): 5222-5226.
31. Samantaray S and Maiti S. "An assessment of genetic fidelity of micropropagated plants of *Chlorophytum borivillianum* using RAPD markers". *Biologia Plantarum* 54.2 (2010): 334-338.
32. Isabel NL., et al. "RAPDs as an aid to evaluate the genetic integrity of somatic embryogenesis derived population of *Picea mariana* (Mill.) B.S.P". *Theoretical and Applied Genetics* 86 (1993): 81-87.
33. Valles MP., et al. "Analysis of genetic stability of plants regenerated from suspension cultures and protoplasts of meadow fescue (*Festuca pratensis* Huds.)". *Plant Cell Report* 12 (1993): 101-106.
34. Mishra RR., et al. "Cyto-morphological and molecular characterization of *Cajanus cajan* × *C. scarabaeoides* F₁ hybrid". *The Nucleus* 55.1 (2012): 27-35.
35. Sahu AR., et al. "In vitro propagation of *Aerva lanata* (L.) Juss through organogenesis". *Indian Journal of Biotechnology* 12.2 (2013): 260.
36. Sahu AR. "In-vitro Propagation of *Aerva lanata* (L.) Juss". *Lambert Academic publication* (2017).
37. Sahu AR. "In-vitro propagation of *Aerva lanata* (L.) Juss through Organogenesis and Somatic embryogenesis" M. Phil. Dissertation, Sambalpur University, Jyoti Vihar, Odisha, INDIA, (2010).

Volume 2 Issue 11 November 2018

© All rights are reserved by Alok Ranjan Sahu and Jogeswar Panigrahi.