



Evaluation of the Potential Use of Seeds and Vegetables in the Formulation of Yogurt

José Villarroel Bastidas^{1*} and Yandry Dayan Zambrano Muñoz²

¹Master's in Food Processing, Research Professor Faculty of Engineering and Production Sciences, Quevedo, Ecuador

²Agroindustrial Engineer, Faculty of Engineering and Production Sciences, Quevedo, Ecuador

*Corresponding Author: José Villarroel Bastidas, Master's in Food Processing, Research Professor Faculty of Engineering and Production Sciences, Quevedo, Ecuador.

Received: March 21, 2022

Published: April 18, 2022

© All rights are reserved by José Villarroel Bastidas and Yandry Dayan Zambrano Muñoz.

Abstract

To take advantage of the nutritional value of chia and canary seed, the technology for making yogurt was researched and developed, which meets the requirements of fermented milk established by INEN standards, by adding seed extracts (chia and canary seed), vegetables (watercress and alfalfa), and two sweeteners (saccharin and stevia). A completely randomized block factorial design A*B*C was applied with three replicates, which corresponds to 24 treatments. To determine the effects between levels, Tukey significance tests ($p < 0.05$) were performed using the Stat Graphics statistical program. The physical-chemical, sensory, and microbiological analyzes carried out among the most important are Protein content with a value of 6.68% in birdseed and 5.24% in chia, while sensory characteristics such as smell, texture and flavor do not present a significant difference, only present in color regarding vegetables, a score of 2.5 out of 5 was obtained in the sensory analyzes in birdseed. The results establish that the best yogurt was the one made from birdseed, enriched with watercress and Stevia because it improves the chemical and sensory characteristics.

Keywords: Alfalfa; Watercress; Sweeteners; Sensory Analysis

Introduction

Yogurt is currently consumed for its healthy benefits and pleasant organoleptic properties, in this way it has become one of the most desired dairy products in the world thanks to the variety of flavors and presentations of different consistencies: liquid, smoothie and semi-solid that exist in the market. market [1]. Agricultural products such as chia, canary seed, watercress, and alfalfa should be of great importance in the daily diet of both children and adults due to their high content of "carbohydrates, proteins, fiber and omega 3" [2].

Yogurt is a dairy product obtained from the fermentation of milk, through the action of suitable microorganisms and result-

ing in a reduction in pH with or without coagulation (isoelectric precipitation) INEN, Ecuadorian Technical Standard [3]. It also provides a balance between nutrients such as: proteins, fats, carbohydrates, vitamins, and minerals necessary for the functioning and maintenance of the body [4].

Hispanic sage, the scientific name for chia, is grown in Santa Elena and Imbabura. Currently, around 1,000 producers of Hispanic salvia [5]. Chia is an annual herbaceous plant that belongs to the *Lamiaceae* family, importance is given to this species because it produces an oil with the highest known content in nature of fatty acid ω -3- α -linolenic, the chia seed has acquired a notorious importance for health and nutrition because its content of omega-3 fatty

acids reduces cardiovascular diseases [6]. It contains about 33% oil with 82.3% essential fatty acids (alpha-linolenic and linoleic acids) [7], between 19 - 23% protein with an adequate profile of essential amino acids and a higher proportion of lysine, methionine and cysteine than in other oilseeds [8].

Canary seed can be used to prepare drinks such as canary seed drink or canary seed milk, which is very beneficial for the body as it contains many nutrients, not like normal milk, but it contains minerals, proteins and vegetable fats which is favorable for it, since it helps to reduce cholesterol [9]. Canary grass contains 18% protein and 8.7% unsaturated lipids; Canary grass extract contains lipase activity and a high nutrient content so it can be an alternative for people who are lactose intolerant or who are on a hypocaloric diet, lipases are carboxylesterases that catalyze the hydrolysis of fatty acids esterified from triacylglycerols containing acid of 10 carbon atoms or more, the amino acids that this grain possesses highlight the unique structure of its proteins, mainly due to its high content of tryptophan [9].

Alfalfa is rich in Vitamins A, B1, B2, B6, C, D, E and K, 10% of its weight are minerals, among which calcium, phosphorus, iron, sulfur, silica, potassium, magnesium stand out. and sodium, 22% of its weight are proteins, whose content increases if the seed has germinated. It also contains isoflavones (*genistein*), coumarins (*coumestrol*), saponins, and digestive enzymes [10].

Watercress is rich in vitamin A (from beta-carotene) and vitamin C, and is a source of folate, calcium, iron, and vitamin E. It also contains useful amounts of vitamin K, thiamin, vitamin B6, potassium, and naturally occurring iodine. It is low in sodium. Due to its high-water content (93%) it is low in calories [11]. It also provides minerals such as calcium, iron and, to a lesser extent, potassium, and phosphorus [12].

Saccharin is considered the oldest sweetener; it was synthesized in early 1879 and has a sweetening power 300 times greater than that of sucrose [13]. The Recommended Daily Intake (RDI) for saccharin is 0-15 mg/kg/day. Stevia is a non-caloric sweetener, it has a large amount of fiber, protein, and minerals, it is a natural sweetener without calories, being 100 to 300 times sweeter than sugar. International organizations endorse its use as a safe supplement [14]. With the widespread appearance of natural sweeteners such as Stevia, which is available in different commercial formula-

tions, the suitability for yogurt needs to be validated with several studies [15].

The research deals with isoelectric precipitation in which a food drink will be obtained, which will be enriched with watercress and alfalfa, sweetened with products such as saccharin and Stevia, thus improving the physicochemical and microbiological characteristics of the product, considering the different formulations. raised.

Materials and Methods

Vegetal material

Seeds: Chia is rich in mucilage, starch and oil; About 2 mm long by 1.5 wide, and oval and glossy, grayish brown to reddish brown, the canary seed the inflorescences are green at first then turn slightly purple, the seeds are shiny brown and sheathed in a small shell. The yogurt-like drink was made with chia seeds imported from Peru and canary seed from Canada.

Vegetables: Alfalfa, due to its great richness in minerals, such as iron, phosphorus, potassium, and calcium, is credited with the ability to strengthen bone mass and, at a preventive level, to be useful for preventing fractures in the elderly and watercress is a plant rich in antioxidants, vitamins and minerals that provide various health benefits, Sweeteners: Stevia and saccharin as natural sweeteners to reduce the content of free sugars in yogurt.

Experimental design

Control yogurt production using skimmed milk placed in the storage tank, pasteurized, and disinfected in a very high temperature (UHT) pasteurization system, adding cultures of *Streptococcus thermophilus Lactobacillus delbrueckii subsp. Bulgaricus*, transfer to the fermentation tanks at a temperature of 43 °C for 4 hours, mix with the different additives and flavorings and finally store at 4 °C.

Yogurts with 50% milk and 50% natural extract (chia and canary seed) were prepared separately, heated for 30 minutes at 80°C, lowered to 43°C, and then inoculated with 1% (vol./vol.) of lactic ferment containing *Streptococcus thermophilus Lactobacillus delbrueckii subsp. bulgaricus* aseptically. The mixes were divided into 2 equal portions; one portion was further inoculated with 8% (vol/vol) watercress and the other with 8% (vol/vol) alfalfa from each probiotic culture. 500 mL of the mixtures were poured into polystyrene glasses and incubated at 43 °C until a pH of 4.5 ± 0.05 was reached. Subsequently, the yogurts were immedi-

ately cooled to 4°C, the sweeteners with Stevia at 1.7% (vol./Vol.) And 1.7 (vol./vol.) Of the saccharin are easily added in the other container, it should be noted that I know Whereas a control was carried out without the addition of any ingredient, the treatments were carried out in triplicate.

Determination of physicochemical parameters

pH measurement

The measurement was carried out at 25 °C (room temperature) using a pH meter, which was calibrated before use with pH 4.0 and pH 7.0 buffers. The AOAC 981.12 method [16] was followed.

Measurement of soluble solids

An electronic refractometer was used, measurements were made at a temperature of approximately 25 °C. This device has a scale from 0 to 100 °Brix. AOAC: Official Methods of Analysis [17].

Determination of titratable acidity

Determination of titratable acidity the titratable acidity (TA), as a percentage of lactic acid, of the yogurt samples was determined according to the Official Methods of Analysis of the Association of Official Agricultural Chemists [18].

$$\%A = (V \times N \times \text{mleq (lactic acid)}) / \text{MW} \times 100 \quad (1)$$

Humidity determination

It was carried out by the gravimetric method Official Methods of Analysis of the Association of Official Agricultural Chemists [18].

Determination of viscosity

The viscosity of the yogurt was determined with a rotational viscometer, model 802 Laboratories, measurements were made at 25 °C, all measurements were recorded at 50 rpm. Yogurt was gently stirred for 20 s (20 s continuous sweeps) before analysis. All determinations were repeated at least 3 times.

Determination of ashes

It was carried out following the AOAC method: Official Methods of Analysis [19], previously drying the samples at 110°C and subsequently calcined at a temperature of 550°C, until constant weight, the results were determined through the following

$$\%C = (W2-W1) / W0 \times 100 \quad (2)$$

Fat determination

The fat content was determined using the Gerber method described by Egan., *et al.* (1987). 9.5 ml of concentrated sulfuric acid, 11 ml of yogurt and 2 ml of fat-free isoamyl alcohol were placed in the butyrometer, the butyrometer was closed and the mixture was stirred for 5 minutes at 1000-1200 rpm. Subsequently, it is placed in a water bath (T = 75-80 °C) for 2 h, after which the fat content is read on the butyrometer scale.

Determination of crude protein

The method is based on the destruction of organic matter with concentrated sulfuric acid, forming ammonium sulfate which, more than sodium hydroxide, releases ammonia, which is distilled into: boric acid, forming ammonium borate, which is titrated with diluted sulfuric acid at a concentration of 0.1 N. It was determined using the Kjeldahl analytical method, which comprises three steps: digestion, distillation, and titration. This protein determination is called crude, because it is not a direct measure of protein, but rather an estimate of total protein based on the nitrogen content of the feed (nitrogen x 6.25 = crude protein) [18].

$$\text{Nitrogen in the sample (\%)} = ((A \times B \times 0.14) / C) \text{ Crude protein (\%)} = \text{Nitrogen in the sample} \times 6.25 \quad (3)$$




Figure 1: Physical chemical analysis BBS = Birdseed, watercress, and saccharin; Control.

Determination of microbiological parameters

E. coli/coliform count

To confirm coliform colonies, a representative number of coliform colonies incubated at 37°C ± 1°C were selected, transferred to a tube containing BGLB broth, and then incubated at 37°C ± 1°C. The number of coliforms per Gram or milliliter was determined by

multiplying the number of confirmed colonies in BGLB by the dilution factor. For confirmation of *E. coli*, a representative number of colonies were selected from plates incubated at 37 °C. Petri film plates can be counted in a standard colony counter or other type of lighted magnifying glass. Incubation time and temperature vary by method. The best-known approved methods are

- AOAC official method 991.14 For coliforms: Incubate 24h ± 2h at 35°C ± 1°C. For *E. coli*: Incubate 48h ± 2h at 35°C ± 1°C.
- AOAC Official Method 998.08 For *E. coli* (meat, poultry, marine): Incubate 24h ± 2h at 35°C ± 1°C.
- NMKL Method (147.1993) For coliforms: Incubate 24h ± 2h at 37°C ± 1°C. For *E. coli*: Incubate 48h ± 2h at 37°C ± 1°C.

Yeasts and molds

Petri film plates are faced in stacks of up to 20 plates at a temperature of 25 °C ± 1 °C for 3-5 days. Read Petri film plates in a standard Quebec-type colony counter or under a magnifying light source. Yeast and mold colonies were numbered separately after 3 and 5 days. Plates were scored according to the number of colonies they contained; cells from the presumably random yeast colonies were examined microscopically to confirm the absence of bacterial colonies.

Sensory evaluation

Consumer acceptance of yogurt was studied after 24hours of storage at 4°C. In a 24-member panel, yogurt was evaluated with samples presented in coded containers in individual cabinets for

sensory analysis, for which they had to rate color, aroma, texture, flavor, and general acceptability on a 5-point hedonic scale: 1 = lowest and 5 = highest.

Statistical treatment

Through the results obtained from a completely randomized design (ANOVA) with a 2*2*2 factorial arrangement for the types of seeds (chia and Canarian seed), vegetables (watercress and alfalfa), and different low-calorie sweeteners (saccharin and stevia. Three repetitions were performed for a total of 24 experimental units for the results of the physical-chemical characterization of the beverage and Tukey's test at the 5% level of significance if there were differences between the combinations. The data were analyzed by the Stat Graphics Centurion XVI program.

Results and Discussion

The characteristics of the yogurt that were developed based on seeds, vegetables and sweeteners are found in the combinations of table 1, the physicochemical analyzes of the yogurt presented representative values in some parameters. It is important to consider that the properties evaluated are not all the parameters that define the quality of yogurt, these parameters are grouped into physical-chemical, microbiological and sensory criteria, which depend on numerous factors and their interactions, and are linked to the main physical components: milk chemicals (fat, protein, and lactose), as well as commonly found micro compounds, such as minerals, vitamins, cholesterol, and terpenes [20].

Physico-chemical parameters

	BAST	BAS	BBS	BBST	CWST	CWS	CAST	CAS	Control
pH	4.48 ^{ABC}	4.66 ^{ABC}	4.64 ^{ABC}	4.61 ^{ABC}	4.70 ^{BC}	4.83 ^C	4.79 ^C	4.71 ^{B^C}	4.44 ^A
Soluble solids	11.33 ^{BC}	12 ^C	11.53 ^{BC}	11.93 ^C	10.53 ^{AB}	9.7 ^A	10.27 ^A	11.5B ^C	16.22 ^D
Acidity	0.77 ^{DE}	0.79 ^E	0.75 ^D	0.71 ^C	0.53 ^A	0.56 ^B	0.57 ^B	0.53 ^A	0.84 ^F
Viscosity	2463.33 ^{AB}	1421.33 ^{AB}	3799.33 ^B	2483A ^B	11440 ^C	17323.33 ^D	14986.67 ^D	17333.33 ^D	1465 ^A
Humidity	79.68 ^B	80.12 ^B	80.02 ^B	79.62 ^B	82.94 ^C	83.03 ^C	83.34 ^C	83.14 ^C	78.32 ^A
Ashes	0.83 ^B	0.8 ^B	0.81 ^B	0.8 ^B	0.96 ^C	0.97 ^C	0.89B ^C	0.88B ^C	0.45 ^A
Protein	6.2 ^D	8.59 ^E	6.3 ^D	5.61 ^{BC}	4.22 ^A	5.31 ^B	5.41 ^B	6.01 ^{CD}	10 ^F
Fat	1.87 ^B	1.57 ^A	1.67 ^A	1.57 ^A	1.97 ^B	1.63 ^A	2.43 ^C	1.67 ^A	2 ^B

Table 1: Physical chemical analysis.

(P < 0.05) BAST: Birdseed, Alfalfa, and Stevia BAS: Birdseed, Alfalfa and Saccharin BBS: Birdseed, watercress and Saccharin BBST: Birdseed, Watercress and Stevia CWST: Chia, Watercress and Stevia CWS: Chia, Watercress and Saccharin CAST: Chia, Alfalfa and Stevia CAS: Chia, Alfalfa and Saccharin, Control.

The pH presented a value of 4.76 in Chia, while Birdseed obtained 4.60, which is higher than the control and the data from [21], in their research on the physicochemical properties and sensory acceptability of fruity goat's milk yogurt. Skimmed with mango and banana. In accelerated tests it is mentioned that the pH ranges between (4.35 - 4.36), the existing variation due to the fact that the pH present in the fruits is lower than the seeds, the pH with a value of 4.73 in Saccharin), while Stevia obtained a value of 4.63, the values are similar to (4.6 - 4.9) of [22], this is due to the enzymatic hydrolysis of lactose into glucose and galactose, glucose is further broken down into lactic acid [23]. This formation of lactic acid causes a decrease in pH.

Soluble solids with a value of 11.70 that corresponds to canary seed, in chia 10.50 were obtained, these data are found below (16 - 24) mentioned [24], in their study on Elaboration of a functional drink of high biological value based on boroj  (Boroj  a patinoi Cuatrec), because the boroj  contains more soluble solids than the seeds. The soluble solids of 11.33 corresponding to Stevia, and saccharin presented a value of 10.88, these values are below the (12.13 - 18.20) reported Physicochemical and flow properties of a settled yogurt enriched with microcapsules containing omega 3 fatty acids [25] in their study of the physicochemical and flow properties of a settled yogurt enriched with microcapsules containing omega 3 fatty acids, the data differ because the sucrose added in the investigation contains more soluble solids than saccharin and stevia and the amounts applied are minimal.

Titrate acidity was observed, a value of 1.17 corresponding to chia, while in Canary seed the value was 0.76, whose data are related to the control and between the parameters of the Andean Standards [26]. ranges from (0.5 - 1.5). In titrate acidity, a value of 0.97 was found in alfalfa, while in watercress it indicated a value of 0.95, the same ones that are between the parameters of (0.6 - 1.2) established by the Salvadoran Standard [26]. The research is within the Codex Alimentarius Standards [4], which indicate that yogurt must have at least 0.6% lactic acid.

Viscosity with a value of 15270.83 in Chia, while the canary seed obtained a value of 2541.75, these values are lower than (5633 - 11717) of [27] the variation in the viscosity of Chia is due to the water absorption capacity in the hydration process that then releases gels that are components of its structure, thus increasing its viscosity, unlike canary grass, the starches they have do not have the facility to retain water and this is reflected in their viscosity.

Humidity, in chia obtained a value of 83.11 and canary seed presented a value of 79.86, the same ones that are between the control ranges and those of [28], in their publication of Fermented Milks in the Community of Madrid, indicating a range of (78.7 - 89.1). The moisture data obtained in saccharin 81.63 and Stevia 81.35, whose values are like (81.06 -87.42) according to [25] in their study of the physicochemical and flow properties of a settled yogurt enriched with microcapsules containing omega 3 fatty acids.

Ash with values of 0.93 in chia, while in Canary seed it obtained a value of 0.81, because the percentages of ash presented by the seeds evaluated as: Canary grass that contains 13.48% in mass production of *Trichoderma harzianum Rifai* in different organic substrates [29], with respect to chia, the content is 4.5%, cited by [30], in his study of the chemical composition of chia, flaxseed and rosehip seeds and its contribution in omega-3 fatty acids, these contents are different from the fruits and milk with which the norm is governed.

Protein of 6.68 that corresponds to Canary seed, in chia it presented 5.24, whose data are higher than (1.86 - 2.84) mentioned [31], in his study on Evaluation of the fermentation of soy beverage with lactic culture, due to the high protein content of the seeds and the concentrations used in the formulas, the same ones found in the parameters of the Official Mexican Standard [32]. The protein with respect to alfalfa with a value of 6.55, while watercress obtained a value of 5.36, these values are like those [33], in their study of rheological changes during the yogurt making process de lupine (*Lupinus mutabilis Sweet*) mentions that the protein obtained from 5.79, both investigations are within the parameters indicated by the Colombian Technical Standard [34] establishes a minimum of 2.6% protein. In proteins, the value of 6.40 in saccharin, while Stevia obtained a value of 5.51, these values are different from those of (3.86 - 3.93) [35], the variation is because sucrose and stevia do not contain protein unlike saccharin.

Fat was found that chia indicated a value of 1.93 and canary seed a value of 1.67, these percentages are within the range established by [4], which indicates that yogurt should have less than 15% fat. In fats, values were found in alfalfa of 1.88 and in watercress a value of 1.71, the same ones that are lower than the Nicaraguan Standards [36], stable a minimum of 3.0% of fat, because those who in this investigation are given, they added different low-fat vegetables, so the percentage varies.

	BAST	BAS	BBS	BBST	CWST	CWS	CAST	CAS	Control
<i>E. coli</i> (UFC/g)	0 ^A	0 ^A	0 ^A	0 ^A	466.67 ^B	333.33 ^B	666.67 ^C	0 ^A	0 ^A
Total coliforms (CFU/g)	0 ^C	0 ^A	0 ^H	300 ^K	500 ^L	100 ^I	100 ^I	0 ^B	0 ^D
Molds/Yeasts (CFU/g)	0 ^H	600 ^I	0 ^F	0 ^G	600 ^K	100 ^I	700 ^L	0 ^C	0 ^A

Table 2: Microbiological analysis.

(P < 0.05) BAST: Birdseed, alfalfa, and Stevia BAS: Birdseed, alfalfa, and saccharin BBS: Birdseed, watercress, and saccharin BBST: Birdseed, watercress, and Stevia CWST: Chia, watercress, and Stevia CWS: Chia, watercress, and saccharin CAST: Chia, alfalfa, and stevia CAS: Chia, alfalfa and saccharin, Control.

In *E. coli* a value of 366.67 was found, which corresponds to chia, in canary seed 0.00 was found, according to the NTE INEN 2395:2011 Standards [37] the ranges (< 1) must be found, the values that conform to the standard they are Canary seed while the chia seed exceeds the parameters due to water retention. In total coliforms, a value of 175.00 was obtained, which corresponds to Chía, while in canary seed the value was 75.00, according to the Standards [37] it is established that the ranges must be between (10 - 100), the value that is adjusted to the canary seed standards while the chia exceeds the parameters. In molds and yeasts, it was found that chia presented a value of 350.00 and canary grass obtained a value of 150.00, which agrees with the parameters established by [37], which indicates a range of (200 - 500).

	BAST	BAS	BBS	BBST	CWST	CWS	CAST	CAS
Colour	1.67 ^A	2.67 ^A	3.33 ^A	3.67 ^A	4 ^A	3 ^A	3 ^A	1.67 ^A
Smell	4.33 ^{AB}	4 ^{AB}	5 ^B	4 ^{AB}	4 ^{AB}	4 ^{AB}	3 ^A	4 ^{AB}
Taste	3 ^A	2.33 ^A	3.33 ^A	2.67 ^A	3.67 ^A	3 ^A	1.67 ^A	2.67 ^A
Texture	1.67 ^{AB}	3.33 ^{ABC}	4 ^{BC}	1.33 ^A	4.67 ^C	3.67 ^{ABC}	3.33 ^{ABC}	3.67 ^{ABC}

Table 3: Sensory analysis.

(P < 0.05) BAST: Birdseed, alfalfa, and Stevia BAS: Birdseed, alfalfa, and saccharin BBS: Birdseed, watercress, and saccharin BBST: Birdseed, watercress, and Stevia CWST: Chia, watercress, and Stevia CWS: Chia, watercress, and saccharin CAST: Chia, alfalfa, and stevia CAS: Chia, alfalfa and saccharin, Control.

The smell presented a value of 4.33 in the canary seed, while the chia obtained a value of 2.67, with respect to the value of the canary seed it is like (4.03 - 4.11) cited by [35], while the chia varies because it has a bean smell that the cupping panel didn't like. A color value of 3.33 was established in watercress, while alfalfa obtained

a value of 2.67, these data are like (2.45 - 3.11) of [38] these values are low because goat milk is added in one of the investigations and the other vegetable extracts therefore the color varied from that of traditional yogurt.

In texture it was found that chia had a value of 3.67 and canary seed a value of 3.33, these values are like (2.20 - 3.38) [23], this is the seeds since goat milk influences the texture being different from that of conventional yogurt. With other investigations on texture, a value of 3.67 was established in saccharin, while stevia obtained a value of 1.67 on a scale of 1 to 5, the sensory responses are like those established [23].




Figure 2: Sensory analysis.

BAST: Birdseed, alfalfa, and Stevia BAS: Birdseed, alfalfa, and saccharin BBS: Birdseed, watercress, and saccharin BBST: Birdseed, watercress, and Stevia CWST: Chia, watercress, and Stevia CWS: Chia, watercress, and saccharin CAST: Chia, alfalfa, and stevia CAS: Chia, alfalfa and saccharin, Control.

Conclusions

The types of chia and canary seeds have a significant direct impact on the protein content of yogurt, in this study it was de-

termined that the canary seed is the seed that most increases the protein content of the drink with a value of 6.68 according to the test of significance (Tukey).

The types of vegetables did not influence the taste, smell, and texture, while there was a difference in color between watercress and alfalfa, therefore, watercress with 3.50 is the most suitable for its application in the yogurt production process.

The types of sweeteners influenced the energy of the final product, therefore, any of the sweeteners (saccharin and Stevia) can be applied, but in small concentrations to keep the calories of the product low unlike traditional sweeteners.

Acknowledgements

To the State Technical University of Quevedo for the support provided in the development of the research.

Financing

It was financed by the Technical University of the State of Quevedo and the Research Directorate that supported the financing of the article.

This work was supported by Dr. Byron Oviedo, former Director of Research, and Dr. Carlos Zambrano, Director of Research.

Contributions of the Authors

Yandry Dayan Zambrano contributed with the development of the experimental phase in which he acquired the different raw materials, in addition to the interpretation of the results of the isoelectric precipitation in which a food drink was obtained, enriched with watercress and alfalfa., sweetened with saccharin and Stevia, José Villarroel Bastidas developed the structure and discussion of the results as well as the conclusions of the combination of the different components such as watercress at 8% (vol/vol) and the other with alfalfa at 8% (vol/ vol) of each probiotic culture, adding the sweeteners 1.7% Stevia (vol/vol) and 1.7 (vol/vol) saccharin.

Data Availability Statement

The yogurt-type drink was made with chia seeds imported from Peru and canary seeds from Canada, while Alfalfa, due to its great richness in minerals, such as iron, phosphorus, potassium and calcium, was obtained in the province of Cotopaxi and Likewise, watercress is a plant rich in antioxidants, vitamins and minerals

that provide various health benefits, sweeteners such as Stevia and saccharin to reduce the content of free sugars in yogurt, obtained in the Quevedo canton, to later elaborate the different combinations in which the incubation time and temperature were incubated in a digital incubation equipment, once the isoelectric precipitation had occurred, the alfalfa was mixed with the watercress, adding the sweeteners Stevia and saccharin. Once the mixtures were finished, the physicochemical, microbiological, and sensory analyzes were carried out.

Bibliography

1. IDNBJA Pinilla. "Estudio Del Proceso De Elaboracion Del Yogurt Batido Con Extracto Natural De Albahaca (Ocimum basilicum L)" (1998): 2-98.
2. RL Zavalía., *et al.* "Desarrollo del cultivo de chía en Tucumán, Republica Argentina". *EEAOC- Advance Agroindustrial* 32 (2010): 27-30.
3. INEN. "NTE INEN 2395 - Leches Fermentadas". *Norma Técnica Ecuatoriana Nte Inen 23952011* (2011): 2-8.
4. Codex Alimentarius. *Leche y Productos Lácteos* (2011).
5. Líderes Revista. "Seis granos andinos se procesan en esta planta" (2018): 2-5.
6. AR Lara., *et al.* "Assessment of the phytochemical and nutritional composition of dark chia seed (*Salvia hispánica* l.)". *Foods* 10.12 (2021).
7. H Busilacchi., *et al.* "EVALUACIÓN DE *Salvia hispánica* l. Cultivada en el sur de Santa fe (República Argentina)". *Cultivation Tropicals* 34.4 (2013): 55-59.
8. RD Victor Aredo., *et al.* "El Método de Superficie de Respuesta y el Modelamiento Difuso en el desarrollo de una galleta con semillas de chía (*Salvia Hispánica* L.) Response". *Agroindustrial Science* (2014).
9. M Cogliatti., *et al.* "Mejoramiento genético de alpiste: Selección y evaluación de líneas de derivadas de la población marroquí PI284184". *Revista de Investigaciones Agropecuarias* 40.2 (2014): 189-195.
10. Inkanat Perú. "Alfalfa: Gran fuente de nutrientes, vitaminas y minerales" (2021).

11. El Telégrafo. "Berro, una planta silvestre con un valor nutricional único".
12. HASN Miguel F, *et al.* "Producción de Berro en Cuautla , Morelos , México Watercress Production in Cuautla , Morelos , Mexico". *Revista Fitotecnica Mexicana* 43 (2020): 543-548.
13. Consumo de edulcorantes no nutritivos: efectos a nivel celular y metabólico". *Perspectivas en Nutrición Humana* 20.2 (2018): 185-202.
14. S Durán A., *et al.* "Estevia (*stevia rebaudiana*), edulcorante natural y no calórico". *Revista Chilena de Nutricion* 39.4 (2012): 203-206.
15. P Narayanan., *et al.* "Use of just-about-right scales and penalty analysis to determine appropriate concentrations of stevia sweeteners for vanilla yogurt". *Journal of Dairy Science* 97.6 (20214): 3262-3272.
16. AOAC Official Method, "AOAC Official Method 981.12". *Control* 8 (2012): 1-2.
17. H Rybak-Chmielewska. "AOAC: Official Methods of Analysis" 1 (2003). doi: 10.7312/seir17116-004.
18. HL Bodily. "Official Methods of Analysis of the Association of Official Agricultural Chemists". *American Journal of Public Health and the Nations Health* 46.7 (1956): 916-916.
19. The Official Methods of Analysis of AOAC International, "AOAC2016". *Official Methods of Analysis AOAC International* 20th Edition (2016).
20. MBLV García., *et al.* "Improvements in goat milk quality: A review". *Small Ruminant Research* (2014): 1-7.
21. V Vásquez-Villalobos., *et al.* "Physicochemical properties and sensory acceptability of goat's milk fruit yogurts with mango and banana using accelerated testing". *Scientia Agropecuaria* 6.3 (2015): 177-189.
22. J Parra Huertas., *et al.* "Comportamiento fisico-quimico de Stevia, Fructosa, Dextrosa y Lactosa como endulzantes a diferentes concentraciones durante el tiempo de incubación en la elaboración de yogurt entero". *Bistua Revista De La Facultad De Ciencias Basicas* 9.2 (2011): 15-20.
23. WN Rojas Castro., *et al.* "Características del yogurt batido de fresa derivadas de diferentes proporciones de leche de vaca y cabra". *Agronomy Mesoamerican* 18.2 (2007): 221.
24. G Salamanca G., *et al.* "Elaboración de una bebida funcional de alto valor biológico a base de Borojo (*Borojoa patinoi* Cuatrec)". *Revista Chilena de Nutrición* 37.1 (2010): 87-96.
25. RC Macedo., *et al.* "Propiedades fisicoquímicas y de flujo de un yogurt asentado enriquecido con microcápsulas que contienen ácidos grasos omega 3". *International Journal of Information Technology* 26.5 (2015): 87-96.
26. N Técnica. "Leches fermentadas. Requisitos" (2007).
27. JA Ruiz Rivera and AO Ramírez Matheus. "Elaboración de yogurt con probióticos (*Bifidobacterium* spp. y *Lactobacillus acidophilus*) e inulina". *Revista de la Facultad de Agronomía* 26.2 (2009): 223-242.
28. P Perez-Landeras. "Leches Fermentadas en la Comunidad de Madrid". *Doc. Técnicos Salud Pública* n° 106 (20166): 1-57.
29. SDI Científica., *et al.* "Producción masiva de *Trichoderma harzianum* Rifai". *Revista Chapingo Serie Horticultura* 14.2 (2008): 185-191.
30. PJPLMSV Quitral. "Composición química de semillas de chía, linaza y rosa mosqueta y su aporte en ácidos grasos omega-3". *Australian Family Physician* 4.2 (2013): 155-160.
31. MC Quicazán., *et al.* "Evaluación de la fermentación de bebida de soya con un cultivo láctico". *Revista Colombiana de Biotecnología* 3.2 (2017): 92-99.
32. SEGOB. "NORMA Oficial Mexicana NOM-181-SCFI-2010". (2010).
33. EVPM Gabriela Segovia. "Cambios reologicos durante el proceso de elaboración de yogurt". *Cienc. e Ing* 16.1 (2007): 123-126.
34. NTC 805. "Productos Lácteos. Leches Fermentadas". *Icontec NTC 805* (2006): 16.

35. P Benjamín Castañeda, *et al.* "Formulación y elaboración preliminar de un yogurt mediante sustitución parcial con harina de tarwi (*Lupinus mutabilis Sweet*)". *Nature Medicine* 3.1 (2009): 2-9.
36. NTN. "Norma tecnica obligatoria nicaragüense" (2007): 1-10.
37. INEN. "Bebida de leche fermentada: requisitos". *Instituto Ecuatoriano de Normalización* (2019): 16-931.

Assets from publication with us

- Prompt Acknowledgement after receiving the article
- Thorough Double blinded peer review
- Rapid Publication
- Issue of Publication Certificate
- High visibility of your Published work

Website: www.actascientific.com/

Submit Article: www.actascientific.com/submission.php

Email us: editor@actascientific.com

Contact us: +91 9182824667