

Volume 6 Issue 6 June 2022

Varieties of USDA Certified Organic *Raphanus Raphanistrum Subsp. Sativus* [L.] Response to Salinity Stress

Jonathan D Hulse^{1*}, Scott Paxon¹, Rachel Meriwether², Laura Tinney¹, John Norvell¹, Joseph Mendrykowski¹, Ralph Streadwick¹, Isabella Leonard¹, Victor Calderas¹, Murrin Reed¹, Anthony Pully¹, Gerald Mahle¹, Katherine Spencer¹, Dominick Gregory¹, Desiree Cao¹, Juliana Rusnak¹, Paiton Jacko³, BayleeAnn Pollett⁴ and Carter West¹

¹Department of Arts and Sciences, Blue Ridge Community College, United States of America

²Department of Horticulture, Blue Ridge Community College, United States of America ³Department of Health Sciences, Blue Ridge Community College, United States of America ⁴Department of Education, Blue Ridge Community College, United States of America

*Corresponding Author: Jonathan D Hulse, Department of Arts and Sciences, Blue Ridge Community College, United States of America. DOI: 10.31080/ASAG.2022.06.1148 Received: April 11, 2022 Published: May 27, 2022 © All rights are reserved by Jonathan D Hulse., *et al.*

Abstract

Raphanus Raphanistrum subsp. Sativus [L.] or radish, is a major economic root crop species grown in the United States of America, and it is a distinct member of the *Brassicaceae* family. It is consumed worldwide for its crunchy texture and piquant flavor, which is often described as sharp or spicy. Globally, agricultural lands have become contaminated with salts in the soils, which can greatly affect productivity of those plots. Climate change is also elevating sea water levels, which has been shown to infiltrate aquifers, and can cause agricultural damage to croplands if used as irrigation water. This experiment tested various sodium chloride concentrations on different varieties of USDA Certified Organic *R. Raphanistrum subsp*. Sativus seeds. The results from this study showed significant statistical differences in germination and root length for different concentrations of sodium chloride. This study also showed significant differences between varieties of *R. raphanistrum subsp*. sativus at different sodium chloride concentrations.

Keywords: Agriculture; Agronomy; Brassicaceae; Raphanus raphanistrum subsp. Sativus; Salinity Stress

Abbreviations

C: Celsius; CMV: Cucumber Mosaic Virus; GMO: Genetically Modified Organism; M: Molar; MiRNA: Micro-RNA; PTES: Potentially Toxic Elements; NaCl: Sodium Chloride; USDA: United States Department of Agriculture; YMV: Youcai Mosaic Virus

Introduction

Brassicaceae is a large and diverse family of plants that includes dozens of economically important species [11]. This family includes the genus *Raphanus*, which is comprised of distinct

species and recently identified subspecies [16,30,43,44]. Of the known species and subspecies in the genus *Raphanus*, some are edible crops, while others are noxious weeds that invade agricultural lands in Europe [43,44]. The cultivated radish belongs to the genus *Raphanus*, but there are discrepancies in species nomenclature [33,11,23,50]. Some research groups refer to cultivated radishes as *Raphanus Raphanistrum* subsp. *Sativus* [L.], while other research groups use *Raphanus sativus* [L.] [1-54]. We will refer to the cultivated radish as *Raphanus Raphanus Raphanistrum* subsp. *Sativus* [L.] for the duration of this study.

Citation: Jonathan D Hulse., *et al.* "Varieties of USDA Certified Organic *Raphanus Raphanistrum Subsp. Sativus* [L.] Response to Salinity Stress". *Acta Scientific Agriculture* 6.6 (2022): 56-64.

Radishes are grown for their roots, leafy greens, seeds, sprouts, and is consumed worldwide for its texture as well as for its unique flavor profile [3,11]. It is understood that the pungent flavor profile of radishes comes from its diverse glucosinolate profile [30]. *R. Raphanistrum* subsp. *Sativus* is grown in Africa, Asia, Europe, North America, South America, and is cultivated in over 35 countries [3]. *R. Raphanistrum* subsp. *Sativus* prefers cool growing conditions and has a fast growth cycle, which makes it ideal for use as a cover crop, or in multiple rotations within agricultural fields [50]. Traditionally, *R. Raphanistrum* subsp. *Sativus* has been grown in soil, but recent research has experimented with alternative soilless growth systems [13,42].

Economically speaking, *R. Raphanistrum* subsp. *Sativus* is a major economic crop, with an annual production of over seven million metric tons, which represents over 2% of all vegetable crops produced worldwide [50]. *R. Raphanistrum* subsp. *Sativus* has many medicinal properties including anti-cancer, anti-microbial, cardio-protective, hepatoprotective, and organoleptic, to name a few [7,19,24,25,33,35,37,50]. *R. Raphanistrum* subsp. *Sativus* is recommended as a food source because of its high content of ascorbate, folate, riboflavin, vitamin A, vitamin B6, and vitamin C [50].

The genome of *R. Raphanistrum* subsp. *Sativus* was sequenced in 2015 by Mitsui., *et al.* and subsequent 'omic' studies have reveled tremendous details into the growth of this economically important organism [27,34,49,52,53]. *R. Raphanistrum* subsp. *Sativus* has been shown to exhibit a 2n = 2x = 18 chromosome profile and has been shown to express over 40 different MiRNA's within their root tissues [51]. Modern 'omic' studies are trying to understand *R. Raphanistrum* subsp. *Sativus*'s response to various stresses, including biotic and abiotic factors [27,34,49,52,53].

Just like any crop species, *R. Raphanistrum* subsp. *Sativus* is susceptible and resistant to a variety of different diseases, which can cause stress to the plant [10,54]. Yoon and colleagues showed that some strains were susceptible to Cucumber Mosaic Virus (CMV), while others were resistant to CMV [54]. In 2017, Choi and colleagues released a first report of Youcai Mosaic Virus (YMV) infecting *R. Raphanistrum* subsp. *Sativus* [10]. Other viral infections of *R. Raphanistrum* subsp. *Sativus* include but are not limited to Papaya Leaf Curl Virus and Pedilanthus Leaf Curl Virus [18,21]. Lee and colleagues report that *R. Raphanistrum* subsp. *Sativus* is susceptible to

the fungal pathogen *Fusarium oxysporum* [Schlecht], while Garibaldi., *et al.* showed susceptibility of radish to *Fusarium equiseti* [Wollenw.] [15,23]. Arefin's research group states that *R. raphanistrum* subsp. *sativus* is susceptible to the fungal pathogen *Alternaria brassicae* [Berk] and *Alternaria brassicicola* [Schwein] [2]. In 2014, Cai and colleagues provided a first report of the fungi *Leptosphaeria biglobosa* [Shoemaker and Brun] on *R. Raphanistrum* subsp. *Sativus* [9]. Petrie (1986) reported *Albugo candida* [Pers.], the causal agent of white rust and stag head, on *R. raphanistrum* subsp. *sativus* [40]. Lakra showed that *R. Raphanistrum* subsp. *Sativus* is susceptible to the downy mildew pathogen *Peronospora parasitica* [Pers.] [22]. Bacterial pathogens of *R. Raphanistrum* subsp. *Sativus* include *Pectobacterium* [Hauben] spp., *Pseudomonas solanacearum* [Smith], *Pseudomonas cannabina pv. Alisalensis* [Gardan], to name a few [26,28].

R. Raphanistrum subsp. *Sativus*'s ability to grow quickly makes it an ideal model organism for agronomic and botanical research. *R. Raphanistrum* subsp. *Sativus* has served as a model organism for a variety of agricultural experiments, including abiotic stress, anti-microbial, biochemical, cover crops, heavy metal accumulation, genetics, and post-harvest spoilage [14,17,29,31,45,47,52]. *R. Raphanistrum* subsp. *Sativus* has been a used as a model crop for developing new techniques for vegetable preservation, which reduces post-harvest waste in food systems [17]. *R. Raphanistrum* subsp. *Sativus* has been a model organism to show associations with diverse insect pollinator species, which is an increasingly important topic in the 21st century [11]. *R. Raphanistrum* subsp. *Sativus* has also been used to shown concentration of Potentially Toxic Elements (PTES) and heavy metals in plant tissues, as a result from irrigation from contaminated waters [4,31].

Utilizing *R. Raphanistrum* subsp. *Sativus* to understand the effects of abiotic stress is increasingly important to researchers as the environment on Earth changes, and food resources become harder to cultivate. Climate change is one of the greatest factors effecting worldwide agricultural production. As the climate changes, abiotic stresses become more prominent, and threaten humans' ability to cultivate food. Extreme weather events and rising ocean waters are a result of climate change. Abiotic stress is often defined as any non-living entity that provides trauma to a host plant, which is not limited to moisture, pH, pollution, salinity, and temperature [6,8,12,47-49,51,52].

Citation: Jonathan D Hulse., et al. "Varieties of USDA Certified Organic Raphanus Raphanistrum Subsp. Sativus [L.] Response to Salinity Stress". Acta Scientific Agriculture 6.6 (2022): 56-64.

It is reported that nearly 50% of agricultural crop lands are affected by salinity due to contaminated irrigation waters [47,48]. When soils become contaminated with salt, plants lose their ability to procure water, and excess salt can negatively affect their metabolism [47,48]. It is reported that *R. Raphanistrum* subsp. *Sativus* has a low to moderate sensitivity to salinity stress [32]. Recent research by Sun., et al. [47,48] and Wang., et al. [51,52] have investigated the molecular mechanisms behind salinity stress in R. Raphanistrum subsp. Sativus. Paromita and colleagues studied the effect of salinity on germination rates of R. Raphanistrum subsp. Sativus cv. Tasakishan Mula-1, Drui, Red Bombay, at 0.66 - 12 ds m⁻¹ of NaCl [39]. Marcelis and Hooijdonk [32] studied the effect of salt stress on R. Raphanistrum subsp. Sativus cv. Saxa Nova, and they investigated the mass of the root after exposure to varying salt concentrations. Norren and Ashraf [38] conducted a study on germination rate of R. Raphanistrum subsp. Sativus cv. Red Neck, Lal Pari, Mino Japani, 40 Days, Mannu Early, and Desi, in different concentrations of sodium chloride (0mM, 60mM, 120mM, 180mM, 240mM). Their study showed a decrease in germination rate as sodium chloride concentrations increased. Overall, there is substantial literature studying the effects of salinity stress on R. Raphanistrum subsp. Sativus, as compared with other major crop species. Unfortunately, little research has investigated USDA Organic varieties of R. Raphanistrum subsp. Sativus, and their ability to tolerate saline conditions.

This study was intended to test the effects of different salinity concentrations on germination rate and root length of two different varieties of USDA Certified Organic *R. Raphanistrum* subsp. *Sativus*. Varying concentrations of sodium chloride (NaCl) were applied to *R. Raphanistrum* subsp. *Sativus* cv. Cherry Belle and *R. Raphanistrum* subsp. *Sativus* cv German Giant seeds in-situ, to test the effects of contaminated irrigation water on the productivity of this economically important species. To date, USDA Certified Organic *R. raphanistrum* subsp. *sativus* cv. Cherry Belle and *R. Raphanistrum* subsp. *Sativus* cv German Giant varieties have not been tested for salinity tolerance in this manner.

Materials and Methods Seed stock

Seeds of *R. Raphanistrum* subsp. *Sativus* cv. Cherry Bell and German Giant were purchased from Back to the Roots[®]. The seeds were marked as USDA Organic, which assures that they are non-GMO, and were certified by ECOCERT SA in Oakland California, USA. *R. Raphanistrum* subsp. *Sativus* cv. Cherry Belle seeds came

from Lot C, while *R. Raphanistrum* subsp. *Sativus* cv. German Giant seeds came from Lot B. The seed packages are labeled as 'Heirloom' varieties, with a sell by date of 12/2022.

Germination rate

Germination rate was measured in-situ by the following technique. Sterile 100 x 15 mm Falcon[®] petri plates were lined with sterile 100 mm medium density Thermo Fisher[®] filter paper disks. Thermo Fisher[®] Transfer pipettes were used to apply three milliliters of Sodium Chloride (NaCl) to the filter paper disks. Concentrations of 0.0 M, 0.1 M, 0.2 M, 0.3 M, 0.4 M, and 0.5 M were used for this experiment. Then, ten seeds of *R. raphanistrum* subsp. *sativus* were evenly placed on the filter paper, and the petri plates were sealed with Parafilm M[®]. The petri plates were placed in natural light under a 14 light/10 dark regiment for one week at 22°C. Replicates of ten petri plates were created for each concentration and each variety of seeds, for a total of 120 petri plates or 1200 seeds. Germination was measured as the percentage of seeds that germinated from 100 seeds. Germination was assessed by the root penetrating the seed coat and emerging onto the filter paper disk.

Measuring root length

Root length was measured after one week of incubation in conditions as mentioned above. Briefly, germinated seeds were removed from the petri plates with sterile forceps and placed on a sterile lab bench. The roots were stretched linearly, and precisely measured in millimeters with a metric ruler. The measurements were recorded in a Microsoft Excel[®] workbook and saved for future analysis.

Statistical analysis

Statistical analysis was completed using Microsoft Excel® - Data Analysis options. The Analysis ToolPak® and Solver Add-In® was installed previous to the statistical analysis. Before ANOVAs, Means, and Two-Tailed T-Tests Assuming Equal Variance were performed in Microsoft Excel®, F-Tests were completed to look for equal variances. Comparisons were made between the control group (0.0M NaCl) to the treatment groups (0.1M, 0.2M, 0.3M, 0.4M, and 0.5M NaCl) for each variety. Average germination rate and average root length was calculated for each treatment and each variety. ANOVAs and T-Tests were used to make comparisons between different concentrations of NaCl and different varieties. Data was recorded into a Microsoft Excel® workbook.

Citation: Jonathan D Hulse., *et al.* "Varieties of USDA Certified Organic *Raphanus Raphanistrum Subsp. Sativus* [L.] Response to Salinity Stress". *Acta Scientific Agriculture* 6.6 (2022): 56-64.

Results and Discussion



Graph 1: Root length of USDA Certified Organic *R. Raphanistrum subsp. Sativus* cv. Cherry Belle and *R. Raphanistrum subsp. Sativus* cv. German Giant under varying concentrations of Sodium Chloride. Error Bars represent an α value of 0.05.



Graph 2: Germination rate of USDA Certified Organic *R. Raphanistrum subsp. Sativus* cv. Cherry Belle and *R. Raphanistrum subsp. Sativus* cv. German Giant under varying concentrations of Sodium Chloride (NaCl). Error Bars represent an α value of 0.05

Average germination rate was calculated for each treatment (0.0M, 0.1M, 0.2M, 0.3M, 0.4M, and 0.5M) NaCl and each variety (Cherry Belle/German Giant) used for this study (Graph 1, graph 2, table 1, table 2). Two-tailed T-Tests (assuming equal variances) were calculated (Table 3) after the data set passed F-Tests. ANVOA's (Single Factor) were calculated to see if there was a difference between the treatment groups used in this study (Supplemental Data). T-Tests were calculated to see if there was a significant difference between germination rate and NaCl concentration, as well

Variety	Concentration (M)	Germination rate (%)	Root Length (mm)
German Giant	0	88	53.9495
German Giant	0.1	64	19.84444444
German Giant	0.2	53	8.35
German Giant	0.3	7	0.365
German Giant	0.4	0	0
German Giant	0.5	0	0

59

Table 1: Average Germination rate and Root length of*R. Raphanistrum* subsp. *Sativus* cv. German Giant.

Variety	Concentration (M)	Germination rate (%)	Root Length (mm)
Cherry Belle	0	85	52.77
Cherry Belle	0.1	79	24.44
Cherry Belle	0.2	46	4.2
Cherry Belle	0.3	7	0.044
Cherry Belle	0.4	6	0.305
Cherry Belle	0.5	0	0

Table 2: Average Germination rate and Root length of USDA

 Certified Organic *R. Raphanistrum* subsp. *Sativus* cv. Cherry Belle.

as root length and NaCl concentration, for both varieties used in this study (Table 3).

Results of the T-Tests showed significant differences between most of the treatment groups (Table 3). *R. Raphanistrum* subsp. *Sativus* cv. Cherry Belle showed significant differences between all treatments except 0.0M vs 0.1M and 0.3M vs 0.4M for germination rate. *R. Raphanistrum* subsp. *Sativus* cv. Cherry Belle showed significant differences between all treatments except 0.3M vs 0.4M, 0.3M vs 0.5M, and 0.4M vs 0.5M for root length. *R. Raphanistrum* subsp. *Sativus* cv. German Giant showed significant differences between all treatments except 0.4M vs 0.5M for germination rate. *R. Raphanistrum* subsp. *Sativus* cv. German Giant showed significant differences between all treatments except 0.3M vs 0.4M, 0.3M vs 0.5M, and 0.4M vs 0.5M for root length.

The data shows that there are significant differences between germination rate and the concentration of NaCl, for both varieties tested during this study. The data also shows that there are significant differences between root length and the concentration of NaCl, for both varieties tested during this study. ANOVA's show

Citation: Jonathan D Hulse., et al. "Varieties of USDA Certified Organic Raphanus Raphanistrum Subsp. Sativus [L.] Response to Salinity Stress". Acta Scientific Agriculture 6.6 (2022): 56-64.

T-Test	CB - GR (%) vs Concentration	CB - RL (mm) vs Concentration	GG - GR (%) vs Concentration	GG - RL (mm) vs Concentration
0 vs 0.1	0.709001756		0.00700526	2.6717E-05
0 vs 0.2	0.006424549	5.18613E-08	2.01093E-05	3.62727E-08
0 vs 0.3	4.178E-10	1.13729E-08	1.53526E-13	1.91419E-09
0 vs 0.4	1.20893E-09	1.25685E-08	5.3254E-16	1.69611E-09
0 vs 0.5	4.53676E-11	1.12485E-08	5.3254E-16	1.69611E-09
0.1 vs 0.2	0.001431412	4.0675E-05	0.008712111	0.008712111
0.1 vs 0.3	1.43288E-11	2.94933E-06	7.76196E-07	5.53675E-05
0.1 vs 0.4	5.59809E-12	9.54016E-06	5.08515E-08	4.41564E-05
0.1 vs 0.5	1.60589E-13	2.87958E-06	5.08515E-08	4.41564E-05
0.2 vs 0.3	0.000157615	0.00087703	3.30229E-07	4.40233E-06
0.2 vs 0.4	0.000103753	0.002085088	6.17274E-09	2.0604E-06
0.2 vs 0.5	1.33723E-05	0.000793039	6.17274E-09	2.0604E-06
0.3 vs 0.4	0.773046143	0.166874593	0.00488525	0.086640135
0.3 vs 0.5	0.015004964	0.040708551	0.00488525	0.086640135
0.4 vs 0.5	0.014234209	0.115248874	#DIV/0!	#DIV/0!

Table 3: Results of the Two-Tailed T-Test (Assuming Equal Variance) for different varietites of *R. Raphanistrum* subsp. Sativus. CB = Cherry Belle, GG = German Giant, GR= Germination Rate, RL = Root Length, Yellow indicates a p-value below the α value of 0.025.

there were significant differences between varieties for root length at 0.2M NaCl. No other differences were observed after ANOVA's and T-Tests were performed.

Overall, R. Raphanistrum subsp. Sativus cv. German Giant had the highest germination rate (88%) for the control (0.0M NaCl), while R. Raphanistrum subsp. Sativus cv. Cherry Belle had a slightly lower germination rate (85%) for the control (0.0M NaCl). At 0.1M NaCl, R. Raphanistrum subsp. Sativus cv. Cherry Belle had a higher germination rate (79%), while R. Raphanistrum subsp. Sativus cv. German Giant had a lower rate of germination (64%). While there is large difference between the germination rate of these two varieties at 0.1M, ANOVA showed a P-vale of 0.08, which is larger than our alpha value of 0.05. With a larger sample size, there might be significant differences between the two varieties at this concentration. At 0.2M NaCl, R. Raphanistrum subsp. Sativus cv. German Giant had the highest germination rate (53%), while R. Raphanistrum subsp. Sativus cv. Cherry Belle had a lower germination rate (46%). At 0.3M NaCl, R. Raphanistrum subsp. Sativus cv. German Giant had a slightly higher germination rate (8%), while R. Raphanistrum subsp. *Sativus* cv. Cherry Belle had a slightly lower germination rate (7%). At 0.4M NaCl, *R. Raphanistrum* subsp. *Sativus* cv. Cherry Belle had a higher germination rate (6%), while none of the *R. Raphanistrum* subsp. *Sativus* cv. German Giant seeds germinated.

The control concentration (0.0M NaCl) showed that *R. Raphanistrum* subsp. *Sativus* cv. Cherry Belle had the largest average root length of 60.66mm, while *R. Raphanistrum* subsp. *Sativus* cv. German Giant had an average root length of 53.94mm. At 0.1M NaCl, *R. Raphanistrum* subsp. *Sativus* cv. Cherry Belle had the largest average root lenth of 24.44mm, while *R. Raphanistrum* subsp. *Sativus* cv. German Giant had an average root length of 19.84mm. Interestingly, at 0.2M NaCl, *R. Raphanistrum* subsp. *Sativus* cv. German Giant had the largest average root length of 8.35mm, while *R. Raphanistrum* subsp. *Sativus* cv. Cherry Belle had an average root length of 4.2mm. At 0.3M NaCl, the root lengths were slightly higher for *R. Raphanistrum* subsp. *Sativus* cv. Cherry Belle had an average average of 0.044mm. At 0.4M NaCl, the root lengths were slightly higher for *R. Raphanistrum* subsp. *Sativus* cv. Cherry Belle 0.305mm,

Citation: Jonathan D Hulse., et al. "Varieties of USDA Certified Organic Raphanus Raphanistrum Subsp. Sativus [L.] Response to Salinity Stress". Acta Scientific Agriculture 6.6 (2022): 56-64.

where the root length for *R. Raphanistrum* subsp. *Sativus* cv. German Giant was 0.0mm. There was no root length data collected for 0.5M, since none of the seeds germinated at that concentration.

This study adds to the knowledge of salinity tolerance in R. Raphanistrum subsp. Sativus, and will hopefully benefit agriculturalists in the future. Previous studies have examined salinity tolerance in R. Raphanistrum subsp. Sativus, but they tested different varieties than this study. Kanjevac and colleagues examined various mechanisms that effected germination rate of R. Raphanistrum subsp. Sativus [20]. Su., et al. examined the effect of different salt concentrations on germination rate and root length of R. Raphanistrum subsp. Sativus, but looked at cv. Jiujin, Changfeng, Shunyuan, Nanpan, May, and Red. [46]. Munir [36] conducted a study to examine the effect of different salt concentrations on root biomass and chlorophyl concentrations of R. Raphanistrum subsp. Sativus cv. Deci White and Lal Pari, but did not examine Cherry Belle or German Giant. Ayyub., et al. [5] investigated the effects of salt stress on the morphological, physiological, and biochemical characters of R. Raphanistrum subsp. Sativus cv. Desi Long, Local Japanese, Laal Pari, 40 Days, Meno, and Green Neck.

Many other studies have examined salinity tolerance, but this expands knowledge to USDA Certified Organic *R. Raphanistrum* subsp. *Sativus* cv. Cherry Belle and *R. Raphanistrum* subsp. *Sativus* cv German Giant. This study will be useful to agriculturalists and farmers who are trying to grow USDA Certified Organic radishes in salt contaminated soils.

Conclusion

Raphanus Raphanistrum subsp. *Sativus* is a major economic crop, and is cultivated world-wide for its roots, leafy greens, seeds, as well as sprouts. Throughout the world, agricultural lands are becoming contaminated with salty ocean water from irrigation with polluted aquifers. *R. Raphanistrum* subsp. *Sativus* was shown to exhibit varying levels of salinity tolerance. The USDA Certified Organic varieties Cherry Belle and German Giant showed different germination rates and different root lengths when exposed to varying concentrations of Sodium Chloride. Differences occurred between varieties at 0.2M NaCl, which is half the salt concentration of the world's oceans. This information will help agriculturalists and farmers find the most adapted variety of radishes for varying levels of salt contamination in agricultural lands.

Acknowledgements

We want to graciously thank Blue Ridge Community College for providing internal funding for the making of this manuscript. No external funding was provided for the creation of this manuscript. Enormous thanks to the many undergraduate researchers who helped collect data for the writing of this manuscript. Many thanks go out to our Chair Laura Baylor Ed.D., Dean Deborah DeWitt Ed.D., and President Laura Leatherwood Ed.D. for their unparalleled leadership at Blue Ridge Community College.

Conflict of Interest: No conflict of interest was committed during the creation of this manuscript.

Bibliography

- Araniti., et al. "The Delay of Raphanus raphanistrum subsp. sativus (L.) Domin Seed Germination Induced by Coumarin Is Mediated by a Lower Ability to Sustain the Energetic Metabolism". Plants 11.7 (2022): 843.
- Arefin M., *et al.* "Integrated use of fungicide, plant extract and bio-agent for management of Alternaria blight disease of Radish (*Raphanus sativus* L.) and quality seed production". *Research in: Agricultural and Veterinary Sciences* 3.1 (2019): 10-21.
- 3. Arro J and JA Labate. "Genetic variation in a radish (*Raphanus sativus* L.) geodiversity collection". *Genetic Resources and Crop Evolution* 69.1 (2022): 163-171.
- 4. Atamaleki Ali., *et al.* "A systematic review and meta-analysis to investigate the correlation vegetable irrigation with wastewater and concentration of potentially toxic elements (PTES): a Case study of spinach (*Spinacia oleracea*) and radish (*Raphanus raphanistrum subsp. sativus*)". *Biological Trace Element Research* 199.2 (2021): 792-799.
- Ayyub CM., et al. "Evaluation of different radish (Raphanus sativus) genotypes under different saline regimes". American Journal of Plant Sciences 7.06 (2016): 894.
- Baek Dongwon., *et al.* "Plant-growth promoting *Bacillus oryzicola* YC7007 modulates stress-response gene expression and provides protection from salt stress". *Frontiers in Plant Science* 10 (2020): 1646.

Citation: Jonathan D Hulse., et al. "Varieties of USDA Certified Organic Raphanus Raphanistrum Subsp. Sativus [L.] Response to Salinity Stress". Acta Scientific Agriculture 6.6 (2022): 56-64.

- Beevi Syed Sultan., *et al.* "Isothiocyanate profile and selective antibacterial activity of root, stem, and leaf extracts derived from *Raphanus sativus* L". *Foodborne Pathogens and Disease* 6.1 (2009): 129-136.
- 8. Behera Tusar Kanti., *et al.* "Approaches Involved in the Vegetable Crops Salt Stress Tolerance Improvement: Present Status and Way Ahead". *Frontiers in Plant Science* 12 (2021).
- Cai X., *et al.* "First report of Leptosphaeria biglobosa causing black leg on *Raphanus sativus* in central China". *Plant Disease* 98.7 (2014): 993-993.
- Choi S-K., et al. "First Report of Youcai mosaic virus in Raphanus raphanistrum subsp. sativus in Korea". Plant Disease 101.11 (2017): 1959.
- Divija SD., et al. "Diversity, abundance and foraging behaviour of insect pollinators in Radish (*Raphanus raphanistrum subsp.* sativus L.)". Journal of Asia-Pacific Entomology (2022): 101909.
- Fan Lianxue., *et al.* "Genome-and Transcriptome-wide characterization of *bZIP* gene family identifies potential members involved in abiotic stress response and anthocyanin biosynthesis in radish (*Raphanus sativus* L.)". *International Journal of Molecular Sciences* 20.24 (2019): 6334.
- Farghly Khaled A., *et al.* "Corn wastes and peanut shell as growing media for production of red radish plants in soilless system". *Communications in Soil Science and Plant Analysis* 51.13 (2020): 1799-1810.
- Ferreira Lucas Borges., *et al.* "Organic carbon and nitrogen contents and their fractions in soils with onion crops in different management systems". *Soil Research* 56.8 (2018): 846-855.
- Garibaldi A., *et al.* "Occurrence of Fusarium equiseti on *Raphanus sativus* seedlings in Italy". *Plant Disease* 101.8 (2017): 1548-1548.
- 16. V Georgescu Mihaela Ioana., *et al. "Raphanus raphanistrum* Subsp. *Landra* (Moretti ex DC.) Bonnier and Layens-Adventitious species of Mediterranean origin adapted as weed in crops-Some considerations on morphological and anatomical peculiarities". *Agriculture and Agricultural Science Procedia* 10 (2016): 123-128.

- Hashemi Seyed., *et al.* "The combined effects of ultrasound and lactic acid in inactivating microorganisms on fresh radish (*Raphanus raphanistrum subsp. sativus*): Microbiological and quality changes". *Food Science and Nutrition* 8.1 (2020): 162-169.
- Ismail H., *et al.* "First report of Pedilanthus leaf curl virus, Tobacco leaf curl betasatellite, and Guar leaf curl alphasatellite infecting radish (Raphanus sativus) in Pakistan". *Plant Disease* 101.5 (2017): 845-845.
- 19. Janardhanan Yashaswini and Mamta Nandakumar Varier. "Isolation and papain digestion of novel antifungal peptides from red radish (Raphanus Raphanistrum subsp sativus) and analysis of selective cytotoxicity for cancer treatment". *Journal of Microbiology, Biotechnology and Food Sciences* 2021 (2021): 611-614.
- 20. Kanjevac Milica., *et al.* "Improvement of Germination and Early Growth of Radish (*Raphanus sativus* L.) through Modulation of Seed Metabolic Processes". *Plants* 11.6 (2022): 757.
- Kumar Susheel., *et al.* "Association of an isolate of papaya leaf curl virus and papaya leaf curl betasatellite with leaf curl disease of radish in India". *Journal of Plant Pathology* 103.2 (2021): 629-634.
- Lakra BS. "Epiphytology and losses of downy mildew (*Peronospora parasitica*) of radish (*Raphanus sativus*) seed crop". Indian Journal of Agricultural Sciences 71.5 (2001): 321-324.
- 23. Lee O New., *et al.* "Genotyping-by-Sequencing-Based Genome-Wide Association Studies of Fusarium Wilt Resistance in Radishes (*Raphanus sativus* L.)". *Genes* 12.6 (2021): 858.
- Lim Hyun-Woo., et al. "Antimicrobial action of Raphanus raphanistrum subsp. sativus (radish) extracts against foodborne bacteria present in various milk products: A Preliminary Study". Journal of Dairy Science and Biotechnology 37.3 (2019): 187-195.
- Lim Hyun-Woo., *et al.* "Organoleptic characteristics of dairy products supplemented with *Raphanus raphanistrum subsp. sativus* (radish) powder: a preliminary study on efficacy against diabetes". *Journal of Dairy Science and Biotechnology* 37.3 (2019): 177-186.

Citation: Jonathan D Hulse., et al. "Varieties of USDA Certified Organic Raphanus Raphanistrum Subsp. Sativus [L.] Response to Salinity Stress". Acta Scientific Agriculture 6.6 (2022): 56-64.

- Lin Chih-Hung., *et al.* "Radish (*Raphanus sativus* L.), a new host of Pseudomonas solanacearum in Taiwan". *Journal of Phytopathology* 3.3 (1994): 147-155.
- 27. Liu Wei., *et al.* "Transcriptome-wide analysis of chromiumstress responsive microRNAs to explore miRNA-mediated regulatory networks in radish (*Raphanus sativus* L.)". *Scientific reports* 5.1 (2015): 1-17.
- Liu H., *et al.* "First report of Pectobacterium carotovorum subsp. brasiliense causing soft rot on *Raphanus sativus* in China". *Plant Disease* 103.6 (2019): 1409-1409.
- 29. Majidi Abdul Hallim. "Effect of different biochar concentration on the growth of three agricultural plants in Afghanistan". *Journal of Wastes and Biomass Management (JWBM)* 4.1 (2022): 01-07.
- Maldini Mariateresa., *et al.* "Identification and quantification of glucosinolates in different tissues of *Raphanus raphanistrum* by liquid chromatography tandem-mass spectrometry". *Journal of Food Composition and Analysis* 61 (2017): 20-27.
- Manzoor Javid., *et al.* "Heavy metals in vegetables and their impact on the nutrient quality of vegetables: A review". *Journal of Plant Nutrition* 41.13 (2018): 1744-1763.
- 32. Marcelis LFM and J Van Hooijdonk. "Effect of salinity on growth, water use and nutrient use in radish (*Raphanus sativus* L.)". *Plant and Soil* 215.1 (1999): 57-64.
- McGehee Cora S., *et al.* "Efficacy of biofungicides against root rot and damping-off of microgreens caused by *Pythium* spp". *Crop Protection* 121 (2019): 96-102.
- 34. Mitsui Yuki., *et al.* "The radish genome and comprehensive gene expression profile of tuberous root formation and development". *Scientific Reports* 5.1 (2015): 1-14.
- 35. Mohammed Ghaidaa Jihadi and Imad Hadi Hameed. "Pharmacological activities: Hepatoprotective, Cardio protective, Anticancer and anti-microbial activity of (*Raphanus raphanistrum subsp. sativus*): A review". *Indian Journal of Public Health Research and Development* 9.3 (2018): 212-217.
- 36. Munir Saba., et al. "Assessment of inter-cultivar variations for salinity tolerance in winter radish (*Raphanus sativus* L.) using photosynthetic attributes as effective selection criteria". World Applied Sciences Journal 21.3 (2013): 384-388.

- Muthusamy Balasubramanian and Girija Shanmugam. "Analysis of flavonoid content, antioxidant, antimicrobial and antibiofilm activity of in vitro hairy root extract of radish (*Raphanus sativus* L.)". *Plant Cell, Tissue and Organ Culture (PCTOC)* 140.3 (2020): 619-633.
- 38. Noreen Zahra and Muhammad Ashraf. "Inter and intra specific variation for salt tolerance in turnip (Brassica rapa L.) and radish (*Raphanus sativus* L.) at the initial growth stages". *Pakistan Journal of Botany* 40.1 (2008): 229-236.
- 39. Paromita Ghosh., *et al.* "Effect of salinity on germination, growth and yield of radish (*Raphanus sativus* L.) varieties". *International Journal of Biosciences (IJB)* 5.1 (2014): 37-48.
- 40. Petrie GA. "Albugo candida on *Raphanus sativus* in Saskatchewan". *Canadian Plant Disease Survey* 66.2 (1986): 43-46.
- Rubio I., *et al.* "First report of the crucifer pathogen Pseudomonas cannabina pv. alisalensis causing bacterial blight on Radish (*Raphanus sativus*) in Germany". *Plant Disease* 96.6 (2012): 904-904.
- 42. Salerno A., *et al.* "Floating system cultivation of radish (*Raphanus sativus* L.): production and quality". *International Symposium on Soilless Culture and Hydroponics* 697 (2004).
- 43. Scott John., et al. "Raphanus raphanistrum L.-wild radish". Biological Control of Weeds in Australia (2012): 486-493.
- 44. Scott John K., *et al.* "Review of the potential for biological control of wild radish (Raphanus raphanistrum) based on surveys in the Mediterranean region". Proceedings of the 13th Australian Weeds Conference. Plant Protection Society of WA, Perth (2002).
- Seo Mi-Suk., *et al.* "Analysis of sugars content by genotypes in 82 radish (*Raphanus sativus* L.)". *Korean Journal of Plant Resources* 31.5 (2018): 453-465.
- Sun Xiaochuan., *et al.* "Identification of novel and salt-responsive miRNAs to explore miRNA-mediated regulatory network of salt stress response in radish (*Raphanus sativus* L.)". *BMC Genomics* 16.1 (2015): 1-16.
- 47. Su Jianjie., *et al.* "Comparison of salt tolerance in Brassicas and some related species". *American Journal of Plant Sciences* 4.10 (2013): 1911.

Citation: Jonathan D Hulse., *et al.* "Varieties of USDA Certified Organic *Raphanus Raphanistrum Subsp. Sativus* [L.] Response to Salinity Stress". *Acta Scientific Agriculture* 6.6 (2022): 56-64.

- Sun Xiaochuan., *et al.* "Unraveling the root proteome changes and its relationship to molecular mechanism underlying salt stress response in radish (*Raphanus sativus* L.)". *Frontiers in Plant Science* 8 (2017): 1192.
- 49. Tang Mingjia., *et al.* "Genome-wide characterization and evolutionary analysis of heat shock transcription factors (HSFs) to reveal their potential role under abiotic stresses in radish (*Raphanus sativus* L.)". *BMC Genomics* 20.1 (2019): 1-13.
- Thuy Luu Thi, *et al.* "Productivity and nutritional quality of radish under different planting geometry". *Journal of Applied Horticulture* 22.2 (2020): 164-168.
- Wang, Jinglei., *et al.* "Bioinformatics analysis of the lipoxygenase gene family in radish (Raphanus sativus) and functional characterization in response to abiotic and biotic stresses". *International Journal of Molecular Sciences* 20.23 (2019): 6095.
- Wang Yan., *et al.* "Genome-wide identification and functional characterization of the cation proton antiporter (CPA) family related to salt stress response in radish (*Raphanus sativus* L.)". *International Journal of Molecular Sciences* 21.21 (2020): 8262.
- Xie Yang., *et al.* "Genome-wide sRNA and mRNA transcriptomic profiling insights into dynamic regulation of taproot thickening in radish (*Raphanus sativus* L.)". *BMC Plant Biology* 20.1 (2020): 1-14.
- Yoon JY., et al. "Evaluation of Radish (Raphanus raphanistrum subsp. sativus L.) Inbred Lines Resistant to Cucumber mosaic virus". The Journal of the Korean Society of International Agriculture (2017).