



## Biorational Approach: An Alternative Approach to Control the Wilt Diseases of Crops

**Ritwik Sahoo\***

MSc, Student, Department of Plant Pathology, Bihar Agricultural University, Sabour, Bhagalpur, Bihar, India

\*Corresponding Author: Ritwik Sahoo, MSc, Student, Department of Plant Pathology, Bihar Agricultural University, Sabour, Bhagalpur, Bihar, India.

DOI: 10.31080/ASAG.2022.06.1181

Received: June 17, 2022

Published: August 30, 2022

© All rights are reserved by Ritwik Sahoo.

### Abstract

Wilt diseases are one of the most severe diseases which vigorously cause losses in various crops like-tomato, brinjal etc. Wilt disease is caused by fungi, bacteria and virus also and can be induced by nematode also. Therefore, wilt disease is a major problem in all over the world. The severity of fusarium wilt in brinjal is 10%-90% [27,36] and for bacterial wilt fruit yield is reduced by 20%-30% [5] in brinjal. Although chemicals are broadly used to control these wilt diseases, but because of growing concerns about health and environmental safety, the use of toxic, carcinogenic, and/or environmentally damaging chemicals are being discouraged. These chemicals leave toxic residues in consumable agricultural commodities. The survey of monitoring of farm-gate samples in different parts of the country recorded pesticide residues above Maximum Residue Limit (MRL) [19,23]. So, novel strategies including biorational approaches against these wilt diseases are needed for their sustainable management. Antagonistic microorganisms, such as fungi and probiotic bacteria from diverse taxonomic genera were found to suppress wilt diseases both *in vitro* and *in vivo*. Various classes of secondary metabolites, such as alkaloids, phenolics, and terpenoids of plant and microbial origin and various plant origin substances significantly inhibit the causal organisms and are also effective in managing the wilt diseases. Besides these organic amendments are also helpful to control these wilt diseases as they are commonly soil borne in nature. Common mode of action of biocontrol agents include-induced resistance, mycoparasitism, antibiosis, antibiotic production, induced systemic resistance and competition. Mechanism of organic amendments include-modification of physico-chemical environment of soil in host favour, pathogen eradication by direct antagonism and decomposition products and microbial metabolites effects. Mode of action of botanicals include-reaction with various enzymes and complex materials in cell wall or membrane. There is proof of control of bacterial wilt by *Pseudomonas sp.* [11] as well green manure derived from medicinal plant [18] and Cinnamon oil [22]. Therefore, these biorational approaches will become very helpful if we use them more commercially than chemicals, this review updates our knowledge in this perspective.

**Keywords:** Wilt; Biorational; Induced Resistance; Mycoparasitism; Antibiosis; Induced Systemic Resistance

### Introduction

The global agricultural production sustains annual loss of about 20 to 30% on an average due to plant diseases on various crops in different countries. Occasionally, the losses rise even to 100% in the most favourable circumstances or when no control measures

are taken in case of some important diseases [16]. Wilt diseases are one of the most severe diseases which vigorously cause losses in various crops like-tomato, brinjal etc. As an example the severity of fusarium wilt in brinjal is 10%-90% [27,36] and for bacterial wilt fruit yield is reduced by 20%-30% [5] in brinjal.

There are various wilt diseases cause severe damages to the crops. It can be due to fungi, bacteria, virus and also can be induced by nematodes. The diseases show various types of symptoms by which we can identify them and go for suitable measures. In Fusarium wilt, the underground stems become dry and brown as a result of cortical decay while roots may have soft and water soaked appearance, stunted growth. One of the first signs of verticillium wilt is the yellowing of older leaves, then yellow changes to brown and then the leaves die. The disease also causes discoloration of vascular tissue, which we may see as brown streaks up the stems of the plants. This discoloration may also be patchy. In bacterial wilt wilting of the foliage is shown followed by collapse of the entire plant. The wilting is characterized by dropping and slight yellowing of leaves and vascular discolouration. Milky white bacterial oozes coming out which is the diagnostic symptom for bacterial wilt when infected stem cut and dipped in water. In case of tomato spotted wilt on approaching the plant, we may notice that the newer foliage has a bronze or purplish tinge to it or it may even die back. Sometimes the plants are stunted or look wilted. Leaf symptoms may first appear as small, light brown flecks. The fruit is usually infected also and may exhibit raised circular areas, sometimes ringed with small brown spots.

Synthetic chemicals may be used in plant protection programs to limit crop damage by pests and pathogens. But because of growing concerns about health and environmental safety, the use of toxic, carcinogenic, and/or environmentally damaging chemicals are being discouraged. These chemicals leave toxic residues in consumable agricultural commodities. The survey of monitoring of farm-gate samples in different parts of the country recorded pesticide residues above Maximum Residue Limit (MRL) [19,23]. That's we should focus on biorational approaches for management of plant diseases. In this review we focus on various biorational approaches to control the wilt diseases of crops.

### Relevance

Wilt diseases of different crops commonly fusarium wilt and bacterial wilt are now a global concern as they severely effect the various crops. Biorational measures can be the alternative measures of it. Though it is a subject of research for decades but has not yet been adequately commercialized. Hence, mechanism and their efficiency should be highlighted as a control measures of wilt diseases. Due to the development of fungicide resistance to

pathogens as well as ecological and public health considerations, the number of available fungicides is reduced day by day. In addition, the industry is facing rising costs of development and the sale of new disease control products. For these reasons, there is a growing need for the development of another disease management products, which can provide a friendly environment and possible economic control of important plant diseases.

### Biorational approach

Biorational agents are designating or relating to pest, disease and weed control agents used because of their specific effects upon the organisms they are intended to control, rather than because of their toxicity or other general effects; employing or involving such agents. These Approaches includes various techniques and agents which are involved in pest and disease control without effecting the other organisms, environment as they are less toxic. As now a days the toxicity of chemicals largely affect the environment, as well as other organisms that are not harmful therefore, we should focus on these approaches.

The advantages of this approach include

- Biorational products can be used to reduce the environmental impact of farming.
- They are less toxic.
- They work in synergy with the plant's microbiome by feeding and/or not killing beneficial microbes.
- They can control the diseases effectively.
- They not only control the pests and diseases but also help the crop to get better nutrient and helps them for better growth [8].

Various biorational approaches to control the wilt theses are-Bio-Control Agents, Organic Amendments, Plant Origin Substances [1,4,9].

### Bio-control agents

Biological Control Agent (BCA) can be defined as the use of natural efficient strains of any microorganisms or modified organisms that reduce the incidence or severity of diseases caused by plant pathogens. It exhibits an antagonistic activity toward a particular phytopathogen [6]. Microbial biocontrol agents (BCAs) for plant diseases are usually fungal or bacterial strains

isolated from the phyllosphere, endosphere or rhizosphere. They are commonly host specific, environment friendly and labor intensive [12].

Various fungal strains for biocontrol are *Trichoderma sp.*, *Gliocladium sp.*, *Aspergillus sp.*, *Fusarium sp.*, *Beauveria sp.* Various bacterial strains are *Pseudomonas sp.*, *Serratia sp.*, *Streptomyces sp.*, *Rhizobium sp.*, *Bacillus sp.*, *Enterobacter sp.* [2,15,20,26,32,37].

**Mechanism of biocontrol**

- Induced Resistance:** Biocontrol agents release various effectors, metabolites, enzymes like callose synthetase which activate the defense reaction of the host by different way like reactive oxygen species, PR proteins-PR1(anti oomycetes, antifungal), PR2(beta-1,3 glucanase), PR3(chitinase), PR4(antifungal), PR6(Proteinase inhibitor). Many compounds are secreted by the *Trichoderma sp.* into the zone of interaction and induce resistance in plants. Fungal proteins such as xylanase, cellulases and swollenins are released by *Trichoderma sp.* [25]. Another group of proteins that induce this mechanism in plants are avirulence-like (Avr) genes [38]. These are produced by a variety of fungal and bacterial plant pathogens. They usually function as race- or pathovar-specific elicitors of hypersensitive and other protection-related responses in plant species that contain the respective resistance (R) gene.
- Mycoparasitism:** Hyper-parasitism is the maximum taken into consideration and the maximum direct form of antagonism. Hyper-parasitism involves tropic increase parasitism of bio manipulate agent toward the goal organism, coiling, very last attack and dissolution of target pathogens cellular wall or membrane with the aid of the activity of enzymes. Harman (2000) reported the involvement of chitinase and  $\beta$ -1,3 glucanase inside the *Trichoderma* mediated biocontrol management. Considering that enzymes are the merchandise of genes, slight alternate in the structure of gene can cause the manufacturing of various enzyme.
- Antibiosis:** by antimicrobial proteins, metabolites like *Trichoderma sp.* produces unique peptaibols, which are peptide antibiotics which can produce pore in the cell membrane and cause programmed cell death. Many microorganisms produce and launch lytic enzymes that may hydrolyze a wide form of polymeric compounds, consisting of chitin,

proteins, cellulose, hemicellulose, and DNA. Expression and secretion of these enzymes by way of specific microbes can now and again result in the suppression of plant pathogen activities immediately. Selvakumar and Srivastava, (2000) reported another antibiosis mechanism is by growth promoting activity. *Aspergillus niger* strain AN-27 was reported to produce growth promoting compounds, 2-carboxy-methyl- 3 -hexyl-maleic anhydride and 2 methyl ene-3- hexyl-butanedioic acid (hexylitaconic acid) that have been immediately answerable for extended root and shoot length and biomass of crop vegetation.

- Competition:** From the microbial perspective, soils and residing plant surfaces are regularly nutrient confined surroundings. So, to colonize the phytosphere, a microbe ought to efficiently compete for the available vitamins [28]. Both the bio control agents and the pathogens compete with one another for the nutrients and space to get established inside the surroundings. Lorito., *et al.* (1994) reported this technique of opposition is taken into consideration to be an oblique interaction among the pathogen and the bio control agent, wherein the pathogens are excluded through the depletion of food base and through physical occupation of the site. Bio controls agents additionally compete with the pathogen for physical occupation of the sites and thereby lessen or postpone the foundation of the pathogen [21].




Figure 1: [21].

Besides these Induced resistance (Production of phytoalexin, PR proteins, lignification), Loon., *et al.* (2000) and Duiiff., *et al.* (1998) reported antibiotic production and competition are also the mechanisms of biocontrol.

**Various fungal Strains with their targeted wilt diseases (Research Achievements)**

Fungal Strains			
Bio-Control Agents	Targeted wilt diseases	Host	References
<i>Trichoderma sp.</i>	Bacterial wilt	Cotton	41
	Fungal wilt	Tomato	20
<i>Gliocladium sp.</i>	Fungal wilt	Tomato	20
<i>Fusarium sp.</i>	Fungal wilt	Tomato	20
<i>Beauveria sp.</i>	Fungal wilt	Capsicum	10

Table 1

**Various bacterial Strains with their targeted wilt diseases (Research Achievements)**

Bacterial Strains			
Bio-Control Agents	Targeted wilt diseases	Host	References
<i>Pseudomonas sp.</i>	Bacterial wilt	Tomato	11
	Fungal wilt	Tomato	32
	Spotted Wilt	Tomato	17
<i>Serratia sp.</i>	Verticillium wilt	Oilseed rape	26
<i>Bacillus sp.</i>	Fungal wilt	Tomato	15
<i>Streptomyces sp.</i>	Fungal wilt	Tomato	3
	Fungal wilt	Tomato	42
	Bacterial Wilt	Tomato	42
<i>Rhizobium sp.</i>	Fungal wilt	Chickpea	31
<i>Enterobacter sp.</i>	Bacterial wilt	Tomato	39

Table 2

**Various Commercial Products of Bioagents**

**Organic amendments**

According to Himanshu., *et al.* 2014) these are the composition of organic moieties usually derived from nature (residues, peat moss from peat bogs), or are the by-products of processing plants

Commercial Products		
Bio-Control Agents	Product	Diseases
<i>Trichoderma sp.</i>	Rootshield	Fungal wilt
<i>Gliocladium sp.</i>	Primastop	Fungal wilt
<i>Fusarium sp.</i>	Fusaclean, Biofox C	Fungal wilt
<i>Bacillus sp.</i>	Biostart	Root affecting Pathogen

Table 3

or mills (sawdust, cedar chips, bark, bagasse, rice hulls) or waste disposal plants (compost, processed sewage sludge, biosolids etc.). Organic Amendments include-Compost, Green Manure, Peat Moss, Fish Meal, Blood Meal, Feather Meal, Oilcakes (Neem, Castor etc.), Wood Chips and Shavings.

**Sources**

According to Nin., *et al.* (2016) and Larney., *et al.* (2012) there are various sources of organic amendments

- **On Farm Sources:** Bedding materials, Compost, Crop residue, Green manure, Silage juice, Weed biomass, Living mulch.
- **Off Farm Sources:** Sphagnum Peat Moss, Wood waste, Non-agricultural wastes (municipal biosolids, food and kitchen waste etc).


**Other sources are**

- **Agriculture Sources:** Livestock manure (fresh, composted), Crop residues, Spent mushroom compost
- **Forestry Sources:** Deinking sludges, Wood chips and Shavings, Wood ash
- **Urban Sources:** Municipal biosolids (sewage sludge, municipal sludge), By-products of food processing industry.

**Mechanism of action**

According to Sharma and Gupta, (2014) various mechanisms include

- **Disease Avoidance:** By modification of physico- chemical environment of soil in host favour
- **Pathogen eradication:** By decomposition products and microbial metabolites effects and through direct antagonism.



**Figure 2:** Possible Pathways of action of organic amendments against pathogen

Source: Chaube and Pundir, 2005.

**Research achievements**

Pathogen	Crop	Organic Amendment	Reference
<i>Fusarium sp.</i>	Several hosts	Vegetal Compost	40
<i>Verticillium Dahliae</i>	Brinjal	Horse manure, Municipal green waste, wood shavings	24
<i>Ralstonia solanacearum</i>	Tomato	Green Manure of the medicinal plant <i>Adhatoda vasica</i> (L.) Nees	18
Tomato Spotted Wilt Virus	Tomato	Alfa alfa straw	7

**Table 4**

**Plant origin substances**

Plant origin substances like-higher plants contain essential oils and a wide spectrum of secondary metabolites such as phenols, flavonoids, quinones, tannins, alkaloids, saponins and sterols. Such plant-derived chemicals may be exploited for their different biological properties. Because of their natural/ plant origin, they are biodegradable and do not usually leave toxic residues or by-products. Commonly used plant substances are get from-Neem (*Azadirachta indica* A. Juss), garlic (*Allium sativum* Linn.),

eucalyptus (*Eucalyptus globulus* Labill.), turmeric (*Curcuma longa* Linn.), tobacco (*Nicotiana tabacum* Linn.) and ginger (*Zingiber officinale* Rosc.) [29].

**Types**

According to Gurjar, *et al.* (2012) there are various types of botanicals -

- **Plant Extracts:** Neem (*Azadirachta indica* A. Juss), garlic (*Allium sativum* Linn.), eucalyptus (*Eucalyptus globulus* Labill.), turmeric (*Curcuma longa* Linn.), tobacco (*Nicotiana tabacum* Linn.) and ginger (*Zingiber offi cinale* Rosc.).
- **Essential Oils:** Nettle oil (*Urtica* spp.), thyme oil (*Thymus vulgaris* Linn.), eucalyptus oil (*Eucalyptus globulus* Labill.), rue oil (*Ruta graveolens* Linn.), lemongrass oil [*Cymbopogon flexuosus* (Steud.) Wats] and tea tree oil (*Melaleuca alternifolia*).
- **Gel and Latex:** *Aloe vera* (Tourn. Ex Linn.)

**Some examples with their mode of action**

Flavonoids-inactivate enzymes, form complex with cell wall, Phenolic acids-bind to adhesions, form complex with cell wall, inactivate enzymes, Phenolics-membrane disruption, substrate deprivation, Terpenoids, essential oils-membrane disruption, Alkaloids-intercalate with cell wall, Tannins-bind to proteins, inhibit enzymes, non-availability of substrate, Coumarins-interaction with eukaryotic DNA, Lectins and polypeptides-form disulphide bridges [29].

**Research achievements**

Pathogen	Botanicals	References
<i>Fusarium sp.</i>	Clove oil, Neem oil, Essential oil of mustard	[8]
<i>Ralstonia solanacearum</i>	Cinnamon oil from <i>Cinnamomum sp.</i>	[22]
Tomato Spotted Wilt Virus	Azadirachtin from Neem Plant	[30]

**Table 5**

**Future prospects**

As additional insights are gained regarding biorational approaches and the mechanisms, there is every reason to believe that the use of biorational approaches will likely control the wilt

diseases of plants and that can also increase food production without imposing any health hazard, and at the same time conserve the environment. It is essential that researchers continue to learn more about biorational approaches and, immediately, translate this knowledge into a form that can readily be used by farmers.

## Conclusion

As chemicals for controlling diseases are toxic to environment, causes health hazards, adversely affect the beneficial microorganisms and if overused it losses its effectivity and as it is not easily degradable so their residues can also affect the same. So now we should focus on various biorational approaches. As we discuss here the effectivity, their mode of action and how they are advantageous than the chemicals. Therefore, we can come to conclusion that we should start the use of these approaches commercially as it will be beneficial for us, our environment and soil ecology. These approaches also indirectly or directly help in plant growth as well as increase the crop yield.

## Bibliography

- Akanmu AO, *et al.* "Plant disease management: leveraging on the plant-microbe-soil interface in the biorational use of organic amendments". *Frontiers in Plant Science* (2021): 1590.
- Aldinary AM, *et al.* "Biocontrol of tomato Fusarium wilt disease by a new Moringa endophytic Aspergillus isolate". *Materials Today: Proceedings* (2021).
- Anitha A and Rabeeth M. "Control of Fusarium wilt of tomato by bioformulation of Streptomyces griseus in green house condition". *African Journal of Basic and Applied Sciences* 1 (2009): 9-14.
- Arora K, *et al.* "Non-edible oil cakes as a novel substrate for DPA production and augmenting biocontrol activity of *Paecilomyces variotii*". *Frontiers in Microbiology* 8 (2017): 753.
- Begum HA. "Studies on the Integrated Management for Tomato Wilt Complex". PhD Thesis, Bangladesh Agricultural University, Mymensingh (2007).
- Beneduzi A, *et al.* "Plant growth-promoting rhizobacteria (PGPR): their potential as antagonists and biocontrol agents". *Genetics and Molecular Biology* 35.4 (2012): 1044-1051.
- Bonanomi G, *et al.* "Organic amendments modulate soil microbiota and reduce virus disease incidence in the TSWV-tomato pathosystem". *Pathogens* 9.5 (2020): 379.
- Bowers JH and Locke JC. "Effect of botanical extracts on the population density of *Fusarium oxysporum* in soil and control of Fusarium wilt in the greenhouse". *Plant Disease* 84.3 (2000): 300-305.
- Cooley DR. "Biorational approaches to disease management in apples". *Biorational Tree-Fruit Pest Management* (2009): 214.
- El Kichaoui A, *et al.* "Development of *beauveria bassiana*-based bio-fungicide against Fusarium Wilt pathogens for Capsicum Annum, a Promising approach toward vital biocontrol industry in Gaza Strip". *IUG Journal of Natural Studies* (2017).
- Frey P, *et al.* "Hrp-mutants of *Pseudomonas solanacearum* as potential biocontrol agents of tomato bacterial wilt". *Applied and Environmental Microbiology* 60.9 (1994): 3175-3181.
- Gerbore J, *et al.* "Biological control of plant pathogens: advantages and limitations seen through the case study of *Pythium oligandrum*". *Environmental Science and Pollution Research* 21.7 (2017): 4847-4860.
- Gupta SURUCHI and Sharma A. "Evaluation of plant yield, macro and micronutrients concentration in spinach (*Spinacia oleracea* L.) plant tissue as well as in soil amended with hair as fertilizer". *International Journal of Chemical Sciences* 12.1 (2014): 73-82.
- Harman GE. "Myths and dogmas of biocontrol changes in perceptions derived from research on *Trichoderma harzianum* T-22". *Plant Disease* 84.4 (2000): 377-393.
- Heidarzadeh N and Baghaee-Ravari S. "Application of *Bacillus pumilus* as a potential biocontrol agent of *Fusarium* wilt of tomato". *Archives of Phytopathology and Plant Protection* 48.13-16 (2015): 841-849.
- Indiaagronet.com, Disease Management (2012).
- Kandan A, *et al.* "Use of *Pseudomonas fluorescens*-based formulations for management of tomato spotted wilt virus (TSWV) and enhanced yield in tomato". *Biocontrol Science and Technology* 15.6 (2005): 553-569.
- Khan RAA, *et al.* "Management of *Ralstonia solanacearum* (Smith) Wilt in Tomato Using Green Manure of the Medicinal Plant *Adhatoda vasica* (L.) Nees". *Gesunde Pflanzen* 72.2 (2020): 129-138.
- Kole RK, *et al.* "Monitoring of pesticide residues in farmgate vegetable samples in West Bengal". *Pesticide Research Journal* 14 (2002): 77-82.

20. Larkin RP and Fravel DR. "Efficacy of various fungal and bacterial biocontrol organisms for control of *Fusarium* wilt of tomato". *Plant Disease* 82.9 (1998): 1022-1028.
21. Latz MA, et al. "Endophytic fungi as biocontrol agents: elucidating mechanisms in disease suppression". *Plant Ecology and Diversity* 11.5-6 (2018): 555-567.
22. Lee YH, et al. "Chemical pesticides and plant essential oils for disease control of tomato bacterial wilt". *The Plant Pathology Journal* 28 (2012): 32-39.
23. Madan VK, et al. "Monitoring of pesticides from farmgate samples of vegetables in Haryana". *Pesticide Research Journal* 8 (1996): 56-60.
24. Malandraki I, et al. "Thermal inactivation of compost suppressiveness implicates possible biological factors in disease management". *Biological Control* 44.2 (2008): 180-187.
25. Martinez C, et al. "Salicylic acid and ethylene pathways are differentially activated in melon cotyledons by active or heat-denatured cellulase from *Trichoderma longibrachiatum*". *Plant Physiology* 127.1 (2001): 334-344.
26. Müller H and Berg G. "Impact of formulation procedures on the effect of the biocontrol agent *Serratia plymuthica* HRO-C48 on *Verticillium* wilt in oilseed rape". *BioControl* 53.6 (2008): 905-916.
27. Nishat S, et al. "Genetic Diversity of the Bacterial Wilt Pathogen *Ralstonia solanacearum* Using a RAPD Marker". *Comptes Rendus Biologies* 338 (2015): 757-767.
28. Pal KK and Gardener BM. "Biological control of plant pathogens (2006).
29. Seethapathy P, et al. "Botanicals in eco-friendly post harvest disease management". *Innovative Farming* 1.3 (2016): 67-71.
30. Shruthi CR, et al. "Bio-efficacy of bio-pesticides, botanicals and new molecules of insecticides against thrips on tomato (2012).
31. Singh S, et al. "Current advancement and future prospect of biosorbents for bioremediation". *Science of the Total Environment* 709 (2020): 135895.
32. Srivastava R, et al. "Evaluation of arbuscular mycorrhizal fungus, fluorescent *Pseudomonas* and *Trichoderma harzianum* formulation against *Fusarium oxysporum* f. sp. *lycopersici* for the management of tomato wilt". *Biological Control* 53.1 (2010): 24-31.
33. Srivastava R, et al. "Evaluation of arbuscular mycorrhizal fungus, fluorescent *Pseudomonas* and *Trichoderma harzianum* formulation against *Fusarium oxysporum* f. sp. *lycopersici* for the management of tomato wilt". *Biological Control* 53.1 (2010): 24-31.
34. Sundaramoorthy S and Balabaskar P. "Biocontrol efficacy of *Trichoderma* spp. against wilt of tomato caused by *Fusarium oxysporum* f. sp. *Lycopersici*". *Journal of Applied Biology and Biotechnology* 1.3 (2013): 0-4.
35. Thambugala KM, et al. "Fungi vs. fungi in biocontrol: An overview of fungal antagonists applied against fungal plant pathogens". *Frontiers in Cellular and Infection Microbiology* 10 (2020): 604923.
36. Vanitha SC, et al. "Bacterial Wilt of Tomato in Karnataka and Its Management by *Pseudomonas fluorescens*". *Biocontrol* 54 (2009): 685-695.
37. Vianene NM and Abawi GS. "*Hirsutella rhossiliensis* and *Verticillium chlamydosporium* as Biocontrol Agents of the Root-knot Nematode *Meloidogyne hapla* on Lettuce". *Journal of Nematology* 32.1 (2000): 85.
38. Woo SL and Lorito M. "Exploiting the interactions between fungal antagonists, pathogens and the plant for biocontrol". In *Novel biotechnologies for biocontrol agent enhancement and management* (2007): 107-130.
39. Xue QY, et al. "Evaluation of the strains of *Acinetobacter* and *Enterobacter* as potential biocontrol agents against *Ralstonia* wilt of tomato". *Biological Control* 48.3 (2009): 252-258.
40. Yogev A, et al. "Plant waste-based composts suppressive to diseases caused by pathogenic *Fusarium oxysporum*". *European Journal of Plant Pathology* 116.4 (2006): 267-278.
41. Yuan S, et al. "Biological control of tobacco bacterial wilt using *Trichoderma harzianum* amended bioorganic fertilizer and the arbuscular mycorrhizal fungi *Glomus mosseae*". *Biological Control* 92 (2016): 164-171.
42. Zheng X, et al. "A *Streptomyces* sp. strain: isolation, identification, and potential as a biocontrol agent against soilborne diseases of tomato plants". *Biological Control* 136 (2019): 104004.