

Rare Case of Trans-Vaginal Evisceration of Small Bowel: Gynecologist's Nightmare

Geetu Bhandoria^{1*}, Nilofer Siddiqui¹ and Sumit Tripathy²¹Department of Obstetrics and Gynecology, Command Hospital, Pune, Maharashtra, India²Department of Obstetrics and Gynecology, Armed Forces Medical College, Pune, Maharashtra, India***Corresponding Author:** Geetu Bhandoria, Department of Obstetrics and Gynecology, Command Hospital, Wanowrie, Pune, Maharashtra, India.**Received:** January 25, 2019; **Publication:** March 14, 2019**Abstract**

Gynecologists are well versed with herniation of viscera (bladder or bowel) through an intact vagina i.e. cystocele or entero-rectocele. Herniation of bowel through a breach in vagina, usually vault is a nightmarish experience. We present a case of Carcinoma vaginal vault who presented with spontaneous evisceration of small bowel.

Keywords: Carcinoma; Vaginal Vault; Evisceration; Colpocleisis; Rectocele

Introduction

Spontaneous evisceration of bowel through vaginal vault is a rare but known complication post hysterectomy. About a hundred case reports have been published in this regard world over [1,2]. There are however even lesser such case reports in cases of gynecological malignancy. This clinical situation is a surgical emergency since viability of bowel is under threat. Also presence or background of malignancy poses additional challenges to management of such cases, as is evident in our case.

Case Report

77 years old postmenopausal lady presented to our outpatient clinic with complaints of mass prolapsing through vaginal vault and bleeding per vaginum of 1 hour duration. She had undergone Total laparoscopic hysterectomy and bilateral salpingo-oophorectomy 7 years back for CIN 3 and Atypical endometrial hyperplasia. She had also undergone Cystocele repair with Sacro-spinous fixation 5 years back for vault prolapse and cystocele. She was found to have Carcinoma vaginal vault 8 months back and is on palliative chemotherapy, in view of advanced disease. On last follow up for carcinoma vault she has developed new peritoneal deposit under umbilicus, suggesting 'progressive disease'.

She gave history of constipation for past one year. On examination, she was apprehensive, afebrile, had slight tachycardia. Local examination revealed small bowel prolapsing through vault. No attempts at vaginal examination or bowel reduction were made in outpatient department. Patient was prepared for emergency exploration and surgery, and shifted to operation theatre.

General anesthesia was administered and both vaginal and laparotomy trolleys were kept ready. Exploration under anesthesia revealed about 4 loops of edematous, though viable, small bowel prolapsing through a rent in vault (Figure 1). There was large rectocele also present, however there was no anterior vaginal or vault descent. GI surgeon confirmed bowel viability and bowel was gently reduced through vault defect after a thorough warm saline wash (Figure 2). 4 cms defect in vault was identified after bowel reposition (Figure 3). There was malignant growth just 2 cms from proximal margin of vault defect. Vaginal wall was thinned out at rent site, also bowel loops were adherent near vault breach site. In view of overall poor prognosis (progressive vault malignancy) and poor vaginal wall integrity a definitive repair was not undertaken. Vault breach was reduced in size with loose interrupted Vicryl 2-0 sutures. Partial Colpocleisis was done, in view of active disease at vault (Figure 4).



Figure 1: Pre-intervention image showing prolapsed and edematous small bowel.