



## Various Classifications of Verrucous Papillary Lesions of Oral Mucosa

**Sachidanand Mallya P\***

Lecturer, Department of Oral Pathology and Microbiology, A B Shetty Memorial Institute of Dental Sciences, Nitte (Deemed to be University), Mangalore

**\*Corresponding Author:** Sachidanand Mallya P, Lecturer, Department of Oral Pathology and Microbiology, A B Shetty Memorial Institute of Dental Sciences, Nitte (Deemed to be University), Mangalore. **E-mail:** mallyapsachin@gmail.com

**Received:** November 20, 2019; **Published:** December 09, 2019

**DOI:** 10.31080/ASDS.2020.04.723

Papillary lesions are those that tunefactive with a cauliflower surface. Some are pedunculated and some are sessile. Some are single, others are multiple or diffusely involve broad area of oral mucosa [1]. Papillary lesions of the oral mucosa are characterized by focal, multifocal or diffuse exophytic growth with cauliflower or verrucous surface irregularity [2].

### Various Classifications are:

#### 1. Based on the type of the lesion [1]

##### a. Reactive/Infectious Lesions

- Squamous Papilloma
- Papillary hyperplasia
- Condyloma Latum
- Condyloma Acuminatum
- Focal Papillary Hyperplasia.

##### b. Neoplasms

- Keratoacanthoma
- Verrucous Carcinoma.

##### c. Idiopathic Lesions

- Pyostomatitis Vegetans
- Verruciform Xanthoma.

#### 2. Based on the number and appearance of lesion (Eversole and Papanicolaou, 1983)

##### a. Focal Papillary and Verrucous Lesions of the Oral Mucosa

- Squamous Papilloma
- Verruca Vulgaris
- Molluscum Contagiosum
- Verruciform Xanthoma

- Sialadenoma Papilliferum
- Keratoacanthoma
- Condyloma Acuminatum
- Squamous Cell Carcinoma
- Warty Dyskeratoma.

##### b. Multifocal papillary and verrucous lesions of the oral mucosa

- Papillary Hyperplasia
- Florid Papillomatosis
- Nevis Unis Lateris
- Verrucous Carcinoma
- Papillary Exophytic Squamous Cell Carcinoma
- Multiple Condylomata
- Focal Epithelial Hyperplasia
- Focal Dermal Hypoplasia Syndrome
- Multiple Hamartoma Syndrome
- Pyostomatitis Vegetans
- Acanthosis Nigricans
- Verruciform Leukoplakia
- Keratosis Follicularis.

#### 3. Based on the involvement of HPV as an etiological factor [2]

- Human papilloma viruses and Head and Neck Lesions: Genotype -Disease
- HPV 2, 4 -Verruca vulgaris
- HPV 6, 11- Condyloma Acuminatum, squamous papilloma
- HPV 13, 32 -Focal epithelial hyperplasia

- HPV 16 -Proliferative verrucous leukoplakia subtypes
- HPV 6, 11, 16 -Verrucous carcinoma
- HPV 16, 18 -Squamous cell carcinoma
- Papillary Oral Lesions without Known Viral Association
- Papillary hyperplasia (Papillomatosis)
- Verruciform Xanthoma
- Cowden syndrome
- Nevus unius lateris
- Acanthosis Nigricans.

**4. Based on their malignant potential [3]**

**a. Benign**

- Viral papillomas
- Squamous papilloma; Verruca vulgaris; Condyloma acuminatum
- Focal epithelial hyperplasia (Heck's disease)
- Fibro-epithelial polyps
- Verruciform xanthoma
- Papillary hyperplasia
- Pyostomatitis vegetans
- Sialadenoma papilliferum
- Acanthosis nigricans
- Darier's disease.

**b. Potentially malignant**

- Verrucous hyperplasia
- Papillary dysplasia
- Proliferative (verrucous) leukoplakia.

**c. Malignant**

- Verrucous carcinoma
- Papillary carcinoma
- Non-invasive (synonymous with papillary dysplasia)
- Invasive (essentially a conventional squamous cell carcinoma requiring treatment as such)
- Carcinoma cuniculatum (essentially a conventional, well differentiated squamous cell carcinoma requiring treatment as such).

**Bibliography**

1. Verrucal-Papillary Lesions. "Textbook of Oral Pathology". PDQ Oral Disease: 140-159.
2. Eversole LR. Human Papillomaviruses and papillary oral Lesions (2000): 144-151.
3. Thomas GJ and Barrett AW. "Papillary and Verrucous Lesions of the oral mucosa". *Diagnostic Histopathology* 15.6 (2009): 279-285.

**Volume 4 Issue 1 January 2020**

© All rights are reserved by Sachidanand Mallya P.