

ACTA SCIENTIFIC GASTROINTESTINAL DISORDERS (ISSN: 2582-1091)

Volume 4 Issue 1 January 2021

Editorial

Hyper-Reactive Malarial Splenomegaly: A Neglected Disease by the Contemporary World

Oumer Abdu Muhie*

Assistant Professor of Medicine, GAMBY Teaching General Hospital, Ethiopia

*Corresponding Author: Oumer Abdu Muhie, Assistant Professor of Medicine, GAMBY Teaching General Hospital, Ethiopia.

Received: November 05, 2020
Published: November 28, 2020

© All rights are reserved by **Oumer Abdu**

Muhie.

Malaria could progress to a chronic form in individuals with partial immunity. Chronic malaria is often characterized by the lack of acute malaria symptoms. Thus, this disorder was considered as asymptomatic carrier state. Hyper-reactive malarial splenomegaly (HMS) is likely the most severe chronic form of malaria. HMS-earlier known as tropical splenomegaly syndrome - has classic manifestations. The manifestations of HMS include, massive splenomegaly, elevated titers of anti-malarial antibodies and polyclonal immunoglobulin M hypergammaglobulinemia [1,2].

HMS diagnosis is challenging as there are various disease which could mimic HMS including visceral leishmaniasis, lymphoproliferative disorders and myeloproliferative disorders. The diagnosis requires exclusion of such other causes and presence of huge splenomegaly usually greater than 10 cm and raised immunoglobulin M.

Data on HMS prevalence is scarce; however reports show its prevalence ranges from 1 - 2% in Nigeria to 80% in some societies in the Papua New Guinea. Though individuals having HMS respond for anti-malarial medications, the syndrome is not correctable. Thus, HMS could relapse after treatment that manifests with reenlargement of the spleen and deterioration of the immunological functions [3,4].

Due to the occurrence of drug resistance, medicines that were effective in the past might not work anymore. Regrettably, no trustworthy records are available and clinical trials are lacking. The response to HMS therapy is assessed by measuring the splenic size, serum IgM titers, hematologic profiles, and by looking at the overall well-being of the patient.

With regards to HMS treatment outcome various studies have shown different outcomes. There are different anti-malarial medications that can be used in the treatment of this disease. Chloroquine, mefloquine, proguanil, pyrimethamine, and the artemisinin derivatives are few of the medications used in the treatment of HMS. The treatment outcome ranged from no change to improvement in almost all the patients. However, these studies have used different regimens and duration of treatment. The response rate ranged from nil to 100 percent [5-8].

HMS has remained a disease with challenging diagnosis, unclear treatment regimen and duration. The World should pay due attention in devising a simple diagnostic technique and formulating an effective treatment modality in tackling this deadly condition.

Bibliography

- Bousema T OL., et al. "Asymptomatic malaria infections: detectability, transmissibility and public health relevance". Nature Reviews Microbiology 12 (2014): 833-840.
- McGregor A DT., et al. "Hyperreactive malarial splenomegaly syndrome-can the diagnostic criteria be improved?" The American Journal of Tropical Medicine and Hygiene 93 (2015): 573-576.
- Mothe B L-CJ., et al. "A case of hyper-reactive malarial splenomegaly". The Role of Rapid Antigen-Detecting and PCR-Based Tests (2007).
- Imelda Bates GB-A. "Review of diagnostic criteria of hyperreactive malarial splenomegaly". The Lancet 349.9059 (1997): 1178.
- Tamayo SG RL., et al. "Examiner dependence on physical diagnostic tests for the detection of splenomegaly: a prospective study with multiple observers". Journal of General Internal Medicine 8.69 (1993): 69-75.

- 6. CR B. "On the clinical detection of enlargement of the spleen". *Australasian Annals of Medicine* 2.78 (1953): 78-80.
- 7. Arkles LB., *et al.* "A palpable spleen is not necessarily enlarged or pathological". *Medical Journal of Australia* 145.15 (1986): 15-17.
- 8. BG Kumar MAS and Chakrapani. "Splenic complications in malaria: a case series". *Southeast Asian Journal of Tropical Medicine and Public Health* 39.5 (2008): 791-794.

Assets from publication with us

- Prompt Acknowledgement after receiving the article
- Thorough Double blinded peer review
- Rapid Publication
- Issue of Publication Certificate
- High visibility of your Published work

Website: www.actascientific.com/

Submit Article: www.actascientific.com/submission.php

Email us: editor@actascientific.com Contact us: +91 9182824667