

## Risk for Newly Diagnosed Diabetes and Impact of Pre-diabetes and Undiagnosed Diabetes on Severity and Mortality for SARS-CoV-2 Infection

Attapon Cheepsattayakorn<sup>1,2\*</sup>, Ruangrong Cheepsattayakorn<sup>3</sup> and Porntep Siriwanarangsun<sup>1</sup>

<sup>1</sup>Faculty of Medicine, Western University, Pathumtani Province, Thailand

<sup>2</sup>10<sup>th</sup> Zonal Tuberculosis and Chest Disease Center, Chiang Mai, Thailand

<sup>3</sup>Department of Pathology, Faculty of Medicine, Chiang Mai University, Chiang Mai, Thailand

\*Corresponding Author: Attapon Cheepsattayakorn, 10<sup>th</sup> Zonal Tuberculosis and Chest Disease Center, Chiang Mai, Thailand.

Received: March 23, 2022

Published: February 13, 2022

© All rights are reserved by Attapon

Cheepsattayakorn., et al.

Hyperglycemia and diabetes are risk factors for critical COVID-19 clinical outcomes [1]. Disproportionately, COVID-19 pandemic has affected increased-risk-of-severe-COVID-19 individuals [2]. During the COVID-19 pandemic, increased in number of type 1 diabetes diagnosed [3,4] and increased frequency and severity of diabetic ketoacidosis (DKA) at the time of diagnosis of diabetes [5] have been demonstrated in European pediatric populations. Independent of diabetic status, pre-diabetes and undiagnosed type 2 diabetes revealed increased risk of severe COVID-19, whereas intrahospital or de novo hyperglycemia predicted critical COVID-19 clinical outcomes [1]. Among individuals aged < 18 years with COVID-19 infection, there was an increased risk for diabetes supported by independent studies in COVID-19 adults [2]. A recent study demonstrated that COVID-19 patients with diabetes presented with more hyperglycemia, whereas type-2-diabetes patients with COVID-19 infection presented with more intensive-care-unit (ICU) need [6]. Another recent study revealed that lower blood glucose concentration, higher inflammatory biomarkers, and ICU admission were associated with diabetes diagnosed at the time of COVID-19 presentation [7].

In conclusion, long-term follow-up studies of COVID-19 are urgently needed to determine the potential relation between COVID-19 and increased risk of diabetes, particularly in persons aged below 18 years.

### Bibliography

1. Vargas-Va'zques A., et al. "Impact of undiagnosed type 2 diabetes and prediabetes on severity and mortality for SARS-CoV-2 infection". *BMJ Open Diabetes Research and Care* 9 (2021): e002026.
2. Barrett CE., et al. "Risk for newly diagnosed diabetes > 30 days after SARS-CoV-2 infection among persons aged < 18 years-United States, March 1, 2020-June 28, 2021". *Morbidity and Mortality Weekly Report* 71.2 (2022): 59-65.
3. Unsworth R., et al. "New-onset type 1 diabetes in children during COVID-19 : multicenter regional findings in the U.K". *Diabetes Care* 43 (2020): e170-e171.
4. Vlad A., et al. "Increased incidence of type 1 diabetes during the COVID-19 pandemic in Romanian children". *Medicina (Kaunas)* 57 (2021): 973.
5. Kamrath C., et al. "Ketoacidosis in children and adolescents with newly diagnosed type 1 diabetes during the COVID-19 pandemic in Germany". *JAMA* 324 (2020): 801-804.
6. Kempegowda P., et al. "Effect of COVID-19 on the clinical course of diabetic ketoacidosis (DKA) in people with type 1 and type 2 diabetes". *Endocrine Connections* 10 (2021): 371-377.
7. Cromer SJ., et al. "Newly diagnosed diabetes vs. pre-existing diabetes upon admission for COVID-19 : associated factors, short-term outcomes, and long-term glycemc phenotypes". *Journal of Diabetes and Its Complications* (2022).

**Assets from publication with us**

- Prompt Acknowledgement after receiving the article
- Thorough Double blinded peer review
- Rapid Publication
- Issue of Publication Certificate
- High visibility of your Published work

**Website:** [www.actascientific.com/](http://www.actascientific.com/)

**Submit Article:** [www.actascientific.com/submission.php](http://www.actascientific.com/submission.php)

**Email us:** [editor@actascientific.com](mailto:editor@actascientific.com)

**Contact us:** +91 9182824667