



## Dietary Habit and Nutritional Status of Regular Students in Jigjiga University, Somali regional state of Ethiopia

**Yimer Mihretie\***

*College of Health Science and Medicine, Department of Human Nutrition, Kotebe Metropolitan University, Ethiopia*

**\*Corresponding Author:** Yimer Mihretie, College of Health Science and Medicine, Department of Human Nutrition, Kotebe Metropolitan University, Ethiopia.

**Received:** February 20, 2019; **Published:** April 01, 2019

### Abstract

**Objective:** To assess the dietary habit and nutritional status of under graduate students in Jigjiga University.

**Methods:** A descriptive survey design was used to answer questions concerning the dietary habit and Nutritional status of regular students in Jigjiga University. Students were interviewed through a structured questioner to assess their nutritional knowledge and dietary habit.

**Results:** A total of 120 respondent students 107(89.2%) of the respondents are age interval of 19-24, 12 (10%) of the respondent are age of 25-29 and at least only 1 (0.8%) student was above 29 old of age. Concerning sex of respondents, 84(70.0%) were male and 36(30%) of respondents were female. Academic year, II and III comprised 120 (100%) of the respondents. From those students 24(20%) were Natural Resource Management (NRM), 24(20%) food science and nutrition, 24(20%) animal range science, 24(20%) crop science and 24(20%) rural development. 70% of students said they were Muslims where by 30% were Christians. Almost 86.7% were singles and at least only 13.3% who are above 25 years old were married. 54.2% engaged themselves in to love relationship JJU Campus. However, 46.8% had no friends in the campus. Most of students by 72.4% believed that love relation in the campus had positive impact on their education.

**Conclusion:** The study showed that the majority of students had normal Body Mass Index (BMI) value and highlighted the presence of unhealthy eating behavior, inadequate nutrient intake of vegetables, snack and high prevalence of under weight among university students.

**Recommendation:** Nutritional education among university students should be encouraged to promote healthier eating habits and life style. There is a need for strategies and coordinated effort at all level (family, university, community) and government to improve the tendency of underweight among university students. Increase the daily fruit (vegetable), snack consumption apart from regular meal would be beneficial for the students.

**Keywords:** Nutritional Status; Dietary Habits; Body Mass Index; Kipping Breakfast; Snack; Fast Food Consumption; Excess Alcohol Consumption.

### Introduction

#### Background

The transition of young people from high school to university has many health implications [1]. It is a time of increased responsibility for food choices and practices. The nutritional knowledge of university students and their dietary habit (food consumption patterns) have received global attention [2]. This because during the transition from secondary school to university, students need to adapt to a new environment. Good eating habits are essential part of a healthy lifestyle [3]. When students fail to adapt adequately this could have negative consequences towards their health habit (behaviors) and subsequent weight status. Eating habit (next to physical activity and sedentary behavior) is an important factor influencing students weight [2]. On the other hand, University stu-

dents tend to engage in problematic eating habit which including unhealthy dieting, high intake of fast food, skipping breakfast, insufficient physical activity, low intake of fruits and vegetables, and minimal consumption of dairy products, although they are generally aware of the negative consequences of those habits.

Rapid changes in physical growth and psychosocial development have placed these young adults as nutritionally vulnerable groups with poor eating habits that fail to meet dietary requirements. Environmental factors also contribute to adoption of unhealthy eating habits among university students [3,4].

Bodyweight and its perception is an important aspect of health and constitute significant role in physical and mental well-being. Generally speaking, evidence related to eating habits or dietary

habit in relation to BMI status in our country especially in university students particularly in AAU which is oldest university in Ethiopia has shown setting is scare. In light of this, it is relevant to investigate and document the prevalence, dietary frequency and physical activity linked to eating habits in relation to BMI difference in Jigjiga University and provides some helpful insight for addressing the issues by the relevant stake holders.

### Statement of Problem

University student's life style may face difficulty in regulating eating behavior since it is the transition of staying away from their parents. Many University Students may be overweight, underweight or obese due to socio demographic and socio economic factors, dietary habits, nutritional knowledge and decrease physical activity. The selection of unhealthy food may have a negative impact on university student's eating behaviors which requires the attention of the university management [5].

It is thus important to deal with the university students' dietary habit (behavior) and uncover the important relevant information in the universities. In line with this, this study was investigate the dietary habit of university students in relation to BMI difference and identifies the determinant factors that are associating with body mass index status among the study participants. However, there is/are no access information which is known about dietary intake and nutritional status of Jigjiga University students which shall be verify and explain in details after this study. Therefore the objective of this study was to assess the dietary habit and nutritional status of under graduate students in Jigjiga University.

### Significant of the Study

There has been no adequate information about standard guideline of good eating practice of food among higher education in Ethiopia. There is a need to map out the magnitude of the problem and avail information within the higher education institute as well as to stakeholders for appropriate intervention. Moreover, the result of the study was also be also helpful for planning and implementing good service provision in higher education as the finding was

indicate which poor nutritional eating habit is more predominant among university students. This research was also served as a base line for researcher who wants to use as a reference and show areas where more research can be done in the future.

## Material and Methodology

### Description of study area

Jigjiga University (JJU) is one of the higher institutions in Ethiopia which has become operational since March, 2007. Jigjiga University is the first and only one university in Somali national regional region state, established in the capital city Jigjiga as direct outcome of government of FDRE policy to expand higher education for all citizens in the different corners of the country to produce complete graduates, conducting problem solving research and provide socially meaningful outer services. Jigjiga University officially started its operation with 712 students and 66 academic and 99 administrative staffs in 3 faculties in the year of 2007. Four years later the student's enrollment has grown to total number of 14,948 students of whom 7804 are regular, 4521 summer, 2623 continue and distance education program Jigjiga on the other hand is located at a distance of 636 km away from Addis-Ababa, eastern part of Ethiopia. Jigjiga lies between 90°2'0" N to 90° 42'0" N and 420°29'00" E to 420°13'00". The mean monthly minimum temperature varies from 5.8°C in November to 14°C from July to September and the mean monthly maximum temperature varies from 25°C in July to 29°C from March to April.

### Sample method

#### The sample size determination

The studies were estimated sample size by employing single population proportion formula. There are 9 faculties in Jigjiga University, this study selected only college of dry land agricultures with all its five departments. Due to their course was block course most of time they have no enough time to use cafeteria properly. Based on the data collected from head of registrar, there are about 1019 students in college of dry land agriculture in all three batches from 2014-2017 GC/2007-2009 EC (Table a).

Data of students of college of dry land Agric from 2014 to 2016, and 2014 to 2017							
No	Departments	Batch of Students					
		Year 2 <sup>nd</sup>			Year 3 <sup>rd</sup>		
		Sex		Total	Sex		Total
		F	M		F	M	
1	Animal and range science	12	21	33	20	49	69
2	Dry land crop science	11	27	38	26	63	89
3	Food science and nutrition	17	18	35	19	71	90
4	Natural resource management	9	27	36	28	70	98
5	Rural development and agricultural extension	25	19	44	28	72	100
	TOTAL			632			

**Table a.** The numbers of male and females students from their respective batches as well as departments.

Study conducted in Ambo university showed that overweight was prevalent in female than males (29.8% versus 24.6%) with total prevalence of 27.5% whereas underweight was more prevalent in males than in females (42.8% % versus 40.7%) with overall prevalence of 40% [6]. Since there is no information regarding nutritional status of students in Jigjiga university we can take consideration from the above mentioned prevalent rate since all University students share many criteria which are the same because of living in the same in all Ethiopian university, under the federal ministry of education. This sample had been calculated by taking the consideration prevalent rate of under nutrition above which is 0.42 at 95% confident sample size had been calculated as:

The sample size had been calculated by using the following formula by Cochran (Israel, 1992), was used to determine the sample size.

$$n = \frac{(Z_{1-\frac{\alpha}{2}})^2 * P * (1 - P)}{d^2}$$

$$[n = \frac{(1.96)^2(0.4)(1-0.4)}{(0.05)^2} = 369 ]$$

369 + 5% non-response rate = 387

Where: n= Total sample size that was be included in this study

- d= marginal error = 0.05
- P= prevalence rate = 0.4
- $Z_{1-\alpha/2}$ = confidence interval (95%) = 1.96

P value of 0.4 was used since there is no information reported before in such study area. Therefore, the final samples included in this study werebeing calculated as follows:

Correction factor:

$$n_c = \left[ \frac{n}{1+(n-1)/N} \right] \dots\dots (1)$$

Where nc - corrected sample size

- n - Initial sample size
- N- Number of source population

$$n_c = \left[ \frac{387}{1+(387-1)/632} \right]$$

$n_c=240$

50% of the correction sample size was considered to check the prevalence of dietary habit and nutritional status of regular students in Jigjiga University

0.50\*240 = 120 sample size

Where:  $n_c$  – is corrected sample size; n –is initial sample size and N-is number of source population.

**Sampling technique**

The sampling technique that has been used is non- probability sampling, in which case respondents meeting the required selection criteria were students and having the criteria mentioned above (all students who registered in 2014 and 2015 as regular

program). They were selected by using convenience sampling. The sampling technique that has been used is non- probability sampling, in which case respondents meeting the required selection criteria were students and having the criteria mentioned above (all students who registered in 2014 and 2015 as regular program). They were selected by using convenience sampling.




Figure 1: Schematic Presentation of the Sampling Technique.

**Data collection method**

**Data collection instruments**

Data had been collected using structured questionnaire prepared in English and was designed by reviewing pertinent research Findings on the issue under caption. The Questionnaire was constituted information on socio demographic, eating attitude, psychological eating behavior and daily meal frequency. The instrument was being pretested in a University which was being selected for the study before the final administration of the questionnaire.

**Anthropometric measures**

- **Weight:** Weight was being measured without shoes and with minimum clothes; to the nearest 100 gm. The scale was be zeroed before each session and calibrated with a known weight. Each participant stood on the scale and the researcher recorded the weight reading.
- **Height:** Height was be measured by using an audiometer, which was be mounted to the wall by resting the body meter on the ground, and placing it against the wall with the visualdisplay facing the researcher and was be measured to the nearest 0.1 cm. The participant was stand straight with heels where weight wasbeing distributed evenly on feet, buttocks and back touching the wall. The head was-being positioned so that the line of vision was at right angles to the body and thearms hang freely by the sides. All measurements werebeing recorded in the questionnaire.

**Study Variable**

**Dependent variable**

- Mental retardness
- Academic weakness
- Health impact

**Independent variable**

- Body mass index (BMI)
- Age
- Sex
- Educational status
- Dietary behavior
- Snacking
- Skipping breakfast
- Alcohol consumption

**Operational Definition of Terms**

- **Underweight:** BMI of male and female students < 18.5 kg/m<sup>2</sup>.
- **Normal weight:** BMI (kg/m<sup>2</sup>) of student's between 18.5 and 24.9 kg/m<sup>2</sup>.
- **Overweight:** BMI of the students > 25.0 kg/m<sup>2</sup> during the period of study.
- **Food Frequency:** Food frequency refers to how often a given food and fluid was usually consumed per day, per week, per month or never.

**Data analysis**

The collected data from each respondent has been entered and analyzed using SPSS version 20 computer Software packages for analysis. Data cleaning was being carried out and cross tabulation was also made for each variables. Odd ratio was also made to assess the strength of associations. The sustainable data was collected and reliable information for dietary habit and nutritional status of regular students in Jigjiga University was obtained

**Ethical considerations**

Ethical clearance and permission was being obtained from research committee in JJU department of food science and nutrition and actual data collection, permission was also obtained from Jigjiga University at Large or in Students Service Director. During the distribution of questionnaires and weight and height measurement taken, students were informed about the purpose and the benefit of the study along with their full right to refuse or completely reject in participation.

**Results and Discussion**

**Results**

Table 1 has shown socio-demographic characters of respondents and related variables among Jigjiga University students in 2017. The finding indicates all students had different bad habits which could affect their learning effectiveness. So, almost half of 53 (44.2%) students were addicted to chewing chat, less than half of them were fanatic for smoking cigarette and using Internet (Facebook), 29 (24.2%) and 28 (23.3%) respectively, and very few 10 (8.3%) students were accustomed to drinking alcohol.

Variable	Frequency	Percent%
1. Age		
2. 19-24	107	89.2
3. 25-29	12	10
4. Other	1	0.8
5. Total	120	100
Sex		
1. Male	84	70.0

2. Female	36	30.0
3. Total	120	100.0
Marital status		
1. Unmarried	104	86.7
2. Married	16	13.3
3. Total	120	100.0
Religion		
1. Orthodox	14	11.9
2. Muslims	94	78.3
3. Protestant	9	7.5
4. Catholic	1	0.8
5. Others	2	1.7
6. Total	120	100.0
Have you a lover in your campus?		
1. Yes	65	54.2
2. No	54	45.0
3. Total	120	100.0
If your answer is yes does your lover disturb your education?		
1. Yes	52	73.2
2. No	18	25.4
3. Total	71	100.0
When do you meet your lover?		
1. When I am free	42	59.2
2. Class time	16	22.5
3. Cafeteria	12	16.9
4. Movie theater	1	1.4
5. Total	120	100.0
From which department you are?		
1. NRM	24	20.0
2. Food science & nutrition	24	20.0
3. Animal & range science	24	20.0
4. Dry land crop science	24	20.0
5. Rural development	24	20.0
6. Total	120	100.0
What is the level of your study year of respondent		
1. 2nd_year	55	45.8
2. 3rd_year	63	52.5
3. Other	2	1.7
4. Total	120	100.0
Have you habit		
1. Yes	32	26.7
2. No	86	71.7
3. Total	120	100.0
If your answer is yes which habit do you have?		
1. Alcohol	10	8.3
2. Smoking	29	24.2
3. Chewing Chat	53	44.2
4. Internet (Facebook)	28	23.3
5. Total	120	100.0

**Table 1.** Socio-demographic character, love and bad habits of students in Jigjiga University, 2017.

Table 2 above indicates that from a total of 120 respondent students, 103(85.8%) respondents have taken meals regularly, and 15(12.5%) have taken meals irregularly. Regarding on the always take breakfast of students, most of the respondents were 102(85%) daily. Regarding on the always take lunch of respondent, most of the students were 102(85%) daily. Regarding on the always take dinner of respondent, most of students were 105(87.5%) daily. Regarding on times eat meals except snacks of respondent, most of the students were 88(73.3%) three times. Regarding on the take snacks apart from regular meals, most of the respondents were 49(40.8%).

Variable	Frequency	Percent (%)
Do you take your meals regularly		
1. Regularly 1000101514637	103	85.8
2. Irregularly	17	14.2
3. Total	120	100.0
Do you always take breakfast?		
1. Daily	102	85.0
2. Three or four times per week	10	8.3
3. Twice per week	4	3.3
4. Rarely	4	3.3
5. Total	120	100.0
Do you always take lunch daily of respondent		
1. Daily	102	85.0
2. Three or four times per week	13	10.8
3. Twice per week	2	1.7
4. Rarely	3	2.5
5. Total	120	100.0
Do you always take dinner daily of respondent?		
1. Daily	105	87.5
2. Three or four times per week	8	6.7
3. Twice per week	2	1.7
4. Rarely	5	4.2
5. Total	120	100.0
How many times do you eat meals except snacks which is one time of respondent?		
1. Once time	13	10.8
2. Two times	13	10.8
3. Three times	88	73.3
4. Four times	6	5.0
5. Total	120	100.0
How often do take snacks apart from regular meals of respondent?		
1. Daily	49	40.8
2. Three or four times per week	37	30.8
3. Once or twice per week	18	15.0
4. Rarely	16	13.3
5. Total	120	100.0

How many meals do you usually eat each day of respondent		
1. One	13	10.8
2. Two	17	14.2
3. Three	83	69.2
4. Four	7	5.8
5. Total	120	100.0
If you skip meals which meals is it usually?		
1. Breakfast	42	35.0
2. Lunch	42	35.0
3. Dinner	36	30.0
4. Total	120	100.0
How often do you eat fruits daily?		
1. Daily	35	29.2
2. Three or four times per week	27	22.5
3. Once or twice per week	23	19.2
4. Rarely	35	29.2
5. Total	120	100.0
How often do you eat fried food?		
1. Daily	41	34.2
2. Three or four times per week	27	22.5
3. Once or twice per week	29	24.2
4. Rarely	23	19.2
5. Total	120	100.0
How often do you take alcohol?		
1. Daily	16	13.3
2. Two or three times per week	16	13.3
3. Rarely	88	73.3
4. Total	120	100.0
How often do you eat with friends and family daily?		
1. Daily	30	25.0
2. Three or four times per week	48	40.0
3. Once or twice per week	20	16.7
4. always alone	22	18.3
5. Total	120	100.0
What type of food do you eat to balance your diet?		
1. Mainly meat	27	22.5
2. Mainly vegetable	25	20.8
3. Meat vegetable and other	47	39.2
4. Variety of food	21	17.5
5. Total	120	100.0
Where do you mostly obtain nutrition information?		
1. Friends	20	16.7
2. Media	32	26.7
3. Parents	31	25.8
4. School	34	28.3
5. Others	3	2.5
6. Total	120	100.0

Table 2. Dietary habits of students (respondents).

Regarding on the how many meals do you usually eat each day, most of the respondents were 83(69.3%) three per day. Regarding on the skip meals, most of the respondents were breakfast 42(35%) and 42(35%) lunch. Regarding on the eat fruits, most of the respondents were 35(29.2%) daily and 35(29.2%) rarely. Regarding on the eat fried, most of the respondents were 41(34.2%) daily. Regarding on the take alcohol, most of the respondents were 88(73.3%) rarely. Regarding on the eat with friends and family, most of the respondents were 48(40%) three or four times per week.

Regarding on the balanced nutrition food, most of the respondents were 47(39.7%) meat, vegetable and other. Regarding on the obtain nutrition information; most of the respondents were 34(28.3%) school. Regarding on the well sleep, most of the respondents were 57(47.5%) hardly sleep well. Regarding on the health condition, most of the respondents were 72(60%) normal. Regarding on the feeling lonely, most of the respondents were 59(49.2%) yes. Regarding on the feel completely out of control when it comes to food, most of the respondents were 73(60.8%) No.

Regarding on the eat until stomach hurts, most of the respondents were 96(80%) No. Regarding on the felt stress, most of the respondents were 68(56.7%) not at all. Regarding on the cigarettes smoke per day, most of the respondents were 114(95%) None. Regarding on the eat because of feeling upset or nervous, most of the respondents were 94(78.3%) No. Regarding on the physical exercise, most of the respondents were 87(72.5%) yes. Regarding on what time physical exercise, most of the respondents were 61(62.2%).

Variable	Frequency	Percent (%)
How well do you sleep?		
1. I hardly ever sleep well	57	47.5
2. I usually do not sleep well	26	21.7
3. Average	37	30.8
4. Total	120	100.0
How is your present health condition?		
1. Bad	5	4.2
2. Normal	72	60.0
3. Good	26	21.7
4. very good	17	14.2
5. Total	120	100.0
Do you eat because of feeling lonely?		
1. Yes	61	50.8
2. No	59	49.2
3. Total	120	100.0

Do you feel completely out of control when it comes to food?		
1. Yes	47	39.2
2. No	73	60.8
3. Total	120	100.0
Do you eat so much until stomach hurts of respondent		
1. Yes	24	20
2. No	96	80.0
3. Total	120	100.0
Have you ever felt stress lately of respondent		
1. Not at all	68	56.7
2. Normal	38	31.7
3. Rather	11	9.2
4. Very much	3	2.5
5. Total	120	100.0
How many cigarettes do you smoke per day of respondent		
1. 1-5	2	1.7
2. 6-10	4	3.3
3. None	114	95.0
4. Total	120	100.0
Do you eat because of feeling upset or nervous of respondent		
1. Yes	11	9.2
2. No	94	78.3
3. Some times	15	12.5
4. Total	120	100.0
Are satisfied by physical exercise of respondent		
1. Yes	87	72.5
2. No	33	27.5
3. Total	120	100.0
At what time you do physical exercise of respondent		
1. Morning	61	62.2
2. Afternoon	39	32.5
3. Others	20	16.7
4. Total	120	100.0

Table 3: Psychological factors affecting students eating habits.

### Discussions

#### Socio-demographic characteristic of students

The sample size of this study was 120. Our finding had revealed that, majority of students by 80% were found in age group of 19-24yrs and at least10% of students were found in 25-29 age group.

Almost 70% of students were male in this study compared to 30% that of female. The same is true sex of students, majority of students in our study are male with 70% compared to 30% of females. 70% of students said they were Muslims where by 30% were Christians. Almost 86.7% were singles and at least only 13.3% who are above 25 years old were marriage. 54.2% engaged themselves in to love relationship JJU Campus. However, 46.8% had no friends in the campus. Most of students by 72.4% believed that love relation in the campus had positive impact on their education.

### Eating habit of students

In this study, breakfast was the most frequently skipped meal among both male and female students. In fact many students in this study skipped breakfast about 35%. However, this finding appeared to be lower when compared with prevalence of skipped breakfast of the other studies such as study conducted in sidamo zone, southern Ethiopia among 11-13yrs age group which was 42.3% [7]. In addis ababa 85.1% (Tefera Tezera, 2017). In another study conducted in Ankara university in turkey reported that, the the percentage of skipping breakfast meals was found to be 82.5% among students (Ergulen S., *et al.* 2001) higher than the percentage of meal taken by students of our study. The main reason for this variation may be lack of nutritional message among JJU students. Our finding here revealed that, almost 87.5% of students in jigjiga university had taken the dinner regularly except snack which seem to be rare among the students. This finding is also in line with study conducted in addis ababa earlier this year where 51.8% of students were reported followed the dining meal regularly (Tefera Tezera, 2017), but there is major different these two studies because majority of JJU students had taken their dinner regularly compared that of addis ababa university in which only portion of student by 51.8%. Taking snack in both finding was said to low however, only 29.1% of student in our study in contrast to 46.9 in addis ababa university students (Tefera Tezera, 2017), the reason for that different may be that those living in capital city spends much of their living outside the campus where they intended to visit all restaurants and cafeteria. majority of student did not consumed alcohol about 91.3% (Tefera Tezera, 2017) and 73% rarely in addis ababa university and jigjiga university respectively. our finding revealed that the above mentioned difference after we compared regular meals, skipped meals, dinner and other relative food eating habit between those studies with our studies may some mentioned reason below: the main reason for this in JJU students is many of students intended to eat outside university cafeteria even if they had no enough money that would satisfied their breast meals and left in dilemma between university cafeteria and outside cafeteria. Eating outside on other hand may expose adolescent from higher income to sedentary way of life style. another for should be lack of awareness about disadvantages of eating disorder so they thought that missing one meals may not have any significant effect to their nutritional status. another reason for this may be attending a university or college students can be stressful experience for

many college students as a result they tend to eat more as a way of getting relief particular in developing countries (Ergulen, Saygamum, 2001), but this may not be the case for Ethiopia.

In Ethiopia weight gain were still considered a sign of healthiness, other shows that overweight as a disease of high economic class [8]. In contrast to this low socio economic status children are at higher risk of becoming obese as compared to those from high socio economic status in developing countries based on the study conducted in Germany (Ergulen, Saygamum, 2001).

The association between eating habits and psychological factors among university students, eating habits score in this study was significantly lower among those who answered 'yes by 49.2%' on the following statements; eat because of feeling lonely 'eat because of feeling bored'. The percentage of students consuming fruit and vegetables one three or four times per week 35% university students. This result is very less as compared to a study conducted in turkey university students where fruit and vegetable consumption was better reported 64.8% of the students [9]. The main reason for that in our finding should be that, there is no availability of fruit and vegetable in jigjiga a city compared to Addis Ababa and other country such as turkey where vegetable and fruits may be available in the market. It had been reported that unhealthy nutritional behaviors such as irregular consumption of meals, low frequency of vegetable and fruit consumption of meals were characteristic of students in general so there is a need to create programs for health education at university students (Uramowski B., *et al.* 2004). Concerning the life style, this study had found that, the sleeping and health condition of the respondents were normal by 60%. In addition to coping with the normal stressors of every day life, university students need to deal with stressors specific to their academics [9]. Although dieting is becoming a popular phenomenon among university students to achieve or maintain a healthy weight, data from this study shows that dieting is not a common practice among the studied population Uramowski B., *et al.* 2004. Likewise, smoking of cigarettes was not common among the students. The prevalence of smoking is low compared to the prevalence of many European countries universities [10]. This finding had revealed that. Alcohol intake was not a common among respondents (jigjiga a university). Nearly half of the students believe that eating meat, vegetables and other food will provide them with balance diet 39.2%. Compared to study of Addis Ababa university students 38% in this study it was agreed that eating a variety of food for good health is important (Tefera Tezera, 2017) [11-50].

### Conclusion

According to the finding of this study, the following conclusions are forwarded. The study showed that the majority of students had normal BMI value and highlighted the presence of unhealthy eating behaviour, inadequate nutrient intake of vegetables, snack and high prevalence of underweight among university students.

Most of the university students skipped breakfast because of fatigue and various reason for instance lack of time and not feeling hungry. Eating behaviours are influence by variety of factors including life style factors. This study showed that eating attitude has an impact on BMI of respondent.

### Bibliography

1. Walid E., *et al.* Relationships between food consumption and living arrangements among university student in four European countries- A cross sectional study 11 (2012): 1-7.
2. Tom Peter C., *et al.* "Determinants of Eating Behavior in University Students: A qualitative study using focus group discussions". *BMC Public Health* 14 (2014): 53.
3. Lua P., *et al.* "Nutrition quality of life among female majority Malay undergraduate students of Health Sciences". 1 (2012): 1-7.
4. Zubaid S., *et al.* "Weight patterns and perceptions among female university students of Karachi: a cross sectional study". *BMC Public Health* 13 (2013): 1-8.
5. Gan W., *et al.* Difference in eating behaviors, dietary intake and body weight status between male and female Malaysia university students 17 (2011): 213-228.
6. Delia S., *et al.* "Self-Reported Sugar- Sweetened Beverage Intake among College Students". 14 (2006): 1825-1831.
7. Anchamo A and Markos B. "Assessment of breakfast eating habits and its association with conitive performance of early adolescents (1113 years) in Shebedino District, Sidama Zone, Southern Ethiopia". *Journal of Food and Nutrition Sciences* 2 (2014): 130-137.
8. Geheber A and Averst A. "Emotional eating in normal weight and underweight individuals". *Eating Behavior* 3 (2003): 341-347.
9. Rakicioglu N and AkalYildiz E. "Energy and nutrient intake and food patterns among Turkish university students". *Nutrition Research and Practice* 5 (2011): 117-123.
10. Science Journal of public Health. eating behaviors of university students in Southern Nigeria: an evaluation of sex difference 2 (2014): 23-27.
11. Aaron P., *et al.* "The fresh man weight gain phenomenon revisited". *Nutrition Reviews* 67.2 (2009): 83-94.
12. Abdallah S., *et al.* "Obesity and eating habits among College Students in Saudi Arabia: A Cross Sectional Study". *Nutrition Journal* 9 (2010): 1-10.
13. Afework M., *et al.* "Nutritional Status of Adolescent Girls from Rural Communities of Tigray, Northern Ethiopia". *Ethiopian Journal of Health Development* 23 (2009): 5-11.
14. Amaia I., *et al.* "Increased cardiovascular risk associated with poor nutritional habits in first -Year university students". *Nutrition Research* 27 (2007): 387-394.
15. Benjamin C. "The Global Epidemic of Obesity: An Overview". *Epidemiology Review* 29 (2007): 1-5.
16. Caille L., *et al.* "Psychosocial and environmental determinants of eating behaviors, physical activity and weight change among college students: A qualitative analysis". *Journal of American College Health* 59 (2011): 531-538.
17. Carvalho H., *et al.* "Risk factors for overweight and obesity in adolescents of Brazilian university: a case-control study". *Nut Hospital* 24 (2009): 17-24.
18. Chapelot D. "The role of snacking in energy balance: A behavioral approach". *The Journal of Nutrition* 141 (2011): 158-162.
19. Cluskey M and Grobe D. "College weight gain and behavior transitions male and female differences". *Journal of the American Dietetic Association* 109 (2009): 325-329.
20. Delva J., *et al.* "The epidemiology of overweight and related lifestyle behaviors- Racial/ethnic and socio economic status differences among American Youth". *American Journal of preventive Medicine* 33 (2007): 178-186.
21. Driskell J., *et al.* "Differences exist in the eating habits of university men and women at fast food restaurants". *Nutrition Research* 26 (2006): 5324-530.
22. Economos C., *et al.* "College freshman stress and weight change: differences by gender". *American Journal of Health Behavior* 32.1 (2008): 16- 25.
23. Greaney M., *et al.* "College student's barriers and enablers for healthful weight management: A Qualitative Study". *Journal of Nutrition Education and Behavior* 41 (2009): 281-286.
24. Gurber K. "Social support for exercise and dietary habits among college students". *Adolescence* 43 (2008): 557-575.
25. Howarth N., *et al.* "Dietary fiber and weight regulation". *Nutrition Reviews* 59 (2001): 129-139.
26. Isganaitis E and Lusting R. "Fast food, central nervous system insulin resistance, and obesity". *Arteriosclerosis, Thrombosis and Vascular Biology* 25 (2005): 2451-2462.
27. Karl P., *et al.* "Prevalence of overweight/obesity and its associated factors among university students from 22 Countries". *International Journal of Environmental Research and Public Health* 11 (2014): 7425-7441.
28. Kathryn S., *et al.* "A Survey of dietary and exercise habits and perceived barriers to following a healthy life style in a college population". 2 (2004): 10-19.



29. Kurubaran G., et al. "Social and Psychological factors affecting eat in habits among university students in a Malaysian medical school: a cross sectional study". *Nutrition Journal* 11 (2012): 1-7.
30. Levitsky D., et al. "The freshman weight gain: A model for the study of the epidemic of obesity". *International Journal of Obesity* 28 (2004): 1435-1442.
31. Meseret Y. "Anthropometric Assessment of Adolescent Malnutrition in Elementary and Secondary Schools of Ambo Town, West Shewa Zone, Oromia Region, Ethiopia". MPH thesis, Addis Ababa University, (2008).
32. Michelle L., et al. "Descriptive epidemiology freshman 15: Fact or Fiction?" 14 (2006): 1438-1443.
33. Natalie P., et al. "Patterns of adolescent physical activity and dietary behaviors". *International Journal Behavioral Nutrition and Physical Activity* 6 (2009): 1-7.
34. Nelosn C., et al. "Emerging adulthood and college aged Youth: An overlooked age for weight- related behavior change". *Obesity* 16 (2008): 2205-2211.
35. Papadaki Scott J and Kapsokefalo U. "Eating habits of university students living at, or away from home in Greece". *Appetite* 49 (2007): 169-176.
36. RouthV, Stern J and Horwithz B. Physiological response of mammals to over nutrition. New -York University Press (1996): 1411-1438.
37. Ruka S., et al. Comparative study of food habits and body shape perception of university students in Japan and Korea (2005).
38. Sakamaki R., et al. "A comparative study of food habits and body shape perceptions of university students in Japan and Korea". *Nutrition Journal* 4 (2005): 1-10.
39. Sallis J and Owen N. "Ecological models of health behaviour". In health behavior and health education: theory, research and practice (2002): 462-484.
40. Sanlie S and Unsan H. "The relationship between body weight and stress and nutritional status in Turkish women". *Pakistan Journal of Nutrition* 6 (2007): 339-344.
41. Sara L., et al. "Electronic feedback in a diet- and physical activity based lifestyle intervention for Weight loss: A randomized controlled trial". *International Journal of Behavior Nutrition and Physical Activity* 8 (2011): 1-8.
42. Savage G., et al. "Snacking behaviors of adolescents and their association with skipping meals". *International Journal of Behavioral Nutrition and Physical Activity* 4 (2007): 1-10.
43. Silliman K., et al. "Survey of dietary and exercise habits and perceived barriers to following a healthy lifestyle in a college population". *Californian Journal of Health Promotion* 2 (2004): 10-19.
44. Story M., et al. "Individual and environmental influences on adolescent eating Behaviors". *Journal of the American Dietetic Association* 102 (2002): 40-51.
45. Strong K., et al. "Weight gain prevention: Identifying theory-based targets for health behavior change in young adults". *Journal of the American Dietetic Association* 108 (2008): 1708-1715.
46. Swinburn B., et al. "Diet, nutrition and the prevention of excess weight gain and obesity". *Public Health Nutrition* 7.1 (2004): 123-146.
47. Walid E., et al. "Correlates of smoking, quit attempts and attitudes towards total smoking ban sat university: Findings from Eleven Faculties in Egypt". *Asian Pacific* 13 (2012): 1-10.
48. World Health Organization. Obesity and overweight September report (2012).
49. World Health Organization. Global health risks: Mortality and burden of disease attributable to selected major risks (2009).
50. Yadav D. Nutritional status of undergraduate students in Pokhara University: An assessment via Body Mass Index 1 (2010): 25-27.

**Volume 3 Issue 4 April 2019**

**© All rights are reserved by Yimer Mihretie.**