



Paediatric Bladder Stones in Kenya. A Single Institution Study

Jumbi T* and Lessan J

Paediatric Surgery Unit, Department of Specialized Surgery, Kenyatta National Hospital, Nairobi, Kenya

***Corresponding Author:** Jumbi T, Paediatric Surgery Unit, Department of Specialized Surgery, Kenyatta National Hospital, Nairobi, Kenya.

Received: September 13, 2021

Published: October 18, 2021

© All rights are reserved by **Jumbi T and Lessan J.**

Abstract

Objective: This study describes the patterns of etiology and presentation of bladder stones in children managed at tertiary facility in Kenya with an aim of highlighting the magnitude of disease.

Materials and Methods: A retrospective review of 20 patients with bladder calculi was done between 2013 and 2020. Data that was collected from the patients' medical records included age and sex distribution; seasonal and geographical distribution; patterns of presentation and etiology.

Results: All patients recruited were male with a mean age of 4.85 years. There was no significant trend in the seasonal and geographical distribution of bladder calculi. Patients with acute urinary retention presented the earliest. Notably, the patients with hematuria in the background of lower urinary tract symptoms presented earlier than those without hematuria. The majority of bladder stones were primary in etiology (n = 12), However 7 patients had secondary bladder stones, resulting from Vesico-ureteric reflux (n = 4), posterior urethral valves (n = 1) and anorectal malformations with recto-urethral fistulas (n = 2).

Conclusion: This study reveals a general inclination towards delayed diagnosis in children presenting with bladder stones especially in the absence of acute urinary retention and hematuria. Additionally, congenital anomalies of the urinary tract should be suspected in patients with bladder stones particularly in presence of urinary tract dilatation.

Keywords: Etiology; Presentation; Bladder Stones; Paediatric; Urolithiasis; Kenya

Abbreviations

LUTS: Lower Urinary Tract Symptoms; KUB: Kidney Ureter Bladder; VCUG: Voiding Cysto-urethrogram; KNH: Kenyatta National Hospital; PUV: Posterior Urethral Valves; ARM: Anorectal Malformation; UTI: Urinary Tract Infection; VUR: Vesico-Ureteral Reflux

Introduction

In developed countries, the prevalence of stone disease is 10 - 15% and bladder calculi account for 5% of the stone burden which

is reported to be higher in developing countries [1]. Bladder stones are more prevalent in males with a reported ratio of 10:1 and peaks at about three years of age [2]. Bladder calculi can be classified as primary, secondary, or migratory [3].

Primary or endemic bladder calculi are defined as those which occur in the absence of other urinary tract pathologies. The determinants for primary bladder stones are nutritional deficiencies especially that of animal protein, poor hydration mainly due to inadequate intake and recurrent diarrhea. Secondary bladder stones

are defined as those which occur in the presence of other urinary tract abnormalities. The determinant for secondary bladder stones include conditions causing functional and mechanical bladder outlet obstruction and conditions causing recurrent urinary tract infections. Migratory bladder stones are defined as those originating from the upper urinary tract [4].

In children bladder stones may cause lower urinary tract symptoms, recurrent infections, pain and hematuria. In the long term, bladder calculi are associated with a risk of damage to the upper tracts and risk of bladder cancer [4]. This study describes the patterns of etiology and presentation of bladder stones in children managed at tertiary facility in Kenya with an aim of highlighting the magnitude of disease.

Materials and Methods

This is a retrospective review of 20 patients managed with bladder calculi between 2013 and 2020 at Kenyatta National Hospital which is the largest teaching and referral hospital in Kenya with a well-established Paediatric surgical unit. Data was collected from the patients’ medical records included age and sex distribution; seasonal and geographical distribution; patterns of presentation and etiology.

All patients were managed according to the hospitals management protocol on bladder calculi which includes clinical assessment; laboratory panel including serum electrolyte measurements and urine analysis; imaging including KUB ultrasound and VCUG; cystoscopy and open cystolithotomy.

Continuous data was analyzed and summarized as mean, median and standard deviations while categorical data was analyzed and displayed in charts by use of frequencies and proportions. Institutional approval to conduct the study was sought and granted.

Results

Age and sex distribution

All the patients recruited were male. The mean age in years was 4.85 while the median age was 5. The youngest patient was 1 year while the oldest patient was 11 years.

Seasonal and geographical distribution

The distribution according to month of presentation is shown in figure 1. The dry seasons in the country are marked in red while

the rainy seasons are marked in blue, there was no seasonal trend noted in this distribution. The highest recorded occurrence was in the month of march with 4 patients.




Figure 1: Seasonal distribution of bladder stone occurrence.

The geographical distribution is shown in figure 2. Most patients reside in central and rift valley of Kenya n = 13 with a few patients distributed in the arid and semi-arid lands n = 7.




Figure 2: Geographical distribution of bladder stone occurrence.

Patterns of presentation

The average duration of symptoms was 1 year. The main presenting complaints included acute urinary retention, hematuria and LUTS such as frequency, urgency, dribbling, strangury and dysuria. Patients with acute urinary retention presented earlier (less than 5 days). Notably, the patients with hematuria in the background of LUTS presented earlier than those without hematuria. This distribution is seen in table 1.

Mode of presentation	Duration of symptoms	Number N = 20	Frequency %
Acute urinary retention	Less than 5 days	6	30%
LUTs with Hematuria	Less than 1 year	4	20%
LUTs without Hematuria	More than 1 year	10	50%

Table 1: Patterns of presentation.

Patterns of etiology

The majority of bladder stones were primary in etiology (n = 12), However 7 patients had secondary bladder stones, the underlying urinary tract pathology in these patients included with Vesicoureteral reflux (n = 4), posterior urethral valves (n = 1) and Anorectal Malformations (ARM) with recto-urethral fistulas (n = 2). This distribution is shown in figure 3.




Figure 3: Patterns of etiology.

VUR- Vesico-Ureteric Reflux.

ARM- Anorectal Malformations.

PUV- Posterior Urethral Valves.

Discussion

Epidemiologic studies have shown an increase in the prevalence pediatric bladder calculi. The true nature of epidemiologic trends may be difficult to study due to under diagnosis and under reporting. Additionally, the detection of risk factors for formation of bladder calculi requires additional investigation to establish the precipitation events in either primary, secondary or migratory stones [1,2]. In this study we highlight the various trends including a propensity of late diagnosis for patients with vague urinary symptomatology and the association of bladder calculi with congenital disorders such as Anorectal malformations, posterior urethral valves and Vesicoureteral Reflux.

In this study all patients presenting with bladder calculi were male. The male to female ratio of bladder calculi is 10:1. The main risk factor for this distribution is the urethral anatomy with girls having short, non-tortuous urethras which allow passage of calculi debris without retaining nuclei which is otherwise retained in the male bladders [2]. The peak age of bladder calculi occurrence is 3 years in children, in this study the average age was 4.85 years however the average duration of symptoms was 1 year making it difficult to conclude on the accurate age of presentation.

Endemicity of bladder stones in Africa has been described previously in relation to the famous afro-asiatic stone-belt. This belt stretches from Morocco, Mauritania, Algeria, Tunisia, Libya, sub-Saharan Africa (Mali, Chad, Rwanda, Ethiopia, Cameroon, Niger), Sudan, Egypt, Turkey, Saudi Arabia, the United Arab Emirates, Iraq, Iran, Afghanistan, Pakistan, India, Nepal, Myanmar, Thailand, and Indonesia to the Philippines [1,5]. Kenya in particular is not included in this belt, however local published literature have noted stone disease in arid and semi-arid lands in Kenya associated with scarcity of drinking water and hot climate with a predisposition to dehydration [6,7]. In this study, a geographical distribution is seen around the central part of Kenya with a few patients scattered in the arid and semi-arid areas. The main catchment for the study setting is central Kenya and may explain the observed distribution. Similarly, the seasonal distribution did not conform to any predisposition of occurrence as there is no much difference in frequency between the wet and dry seasons of the year. This has been documented in previous local literature where the distribution of urolithiasis was evenly distributed throughout the seasons in a six-year period [7].

The presentation of bladder calculi includes presence of Lower Urinary tract symptoms such as frequency, urgency, dribbling, strangury and dysuria. In addition, there may be reports of hematuria, turbid sandy urine or passing of actual stones per urethra. If complete obstruction ensues the presentation will be that of acute urinary retention, this may be accompanied by other symptomatology of straining including rectal prolapse and conjunctival hemorrhages. Finally, bladder calculi may only present with recurrent UTI and may be an incidental finding in upto 10% of patients [1,5,8]. This study revealed an interesting pattern of presentation with an average duration of symptoms of upto 1 year. This delay in presentation was more in patients without hematuria and without complete urine retention. On the other hand, patients with LUTs and hematuria presented much earlier. Logically, the patients with acute urinary retention presented the earliest. This pattern conforms to the poor health seeking behavior seen in developing countries, in particular a local study carried out in Kenya shows that caregivers who delay in taking their children to the hospital immediately after symptomatology usually perceive the illness as mild and self-limiting [9].

The majority of primary/endemic stones notwithstanding, the occurrence of secondary bladder stones and their pattern of distribution in this study is worth discussing. Secondary bladder stones contributed to almost one third of calculi. Majority of these had Vesico-ureteric reflux (VUR), a condition which is documented to predispose to stone formation through several mechanisms. VUR is linked to urinary stasis, bacteraemia and recurrent urinary tract infections which are known to predispose to stone formation [10]. Secondly, studies confirm that there is an association between hypercalciuria and hyperuricosuria with VUR which leads to urinary stone formation [10,11].

The association of Posterior Urethral Valves and vesical calculi has been reported in various published case reports. The highlighted objective in these reports is to have clinical awareness of the existence of such an association which may otherwise lead to delayed diagnosis of posterior urethral valves [12,13]. In one of our patients, this association was detected and the patient was treated by both valve ablation and open cystolithotomy.

A rare but possible association is the presence of bladder calculi in patients with anorectal malformations [14]. ARM is a common paediatric surgical condition that presents with recto-urinary fistu-

las such as recto-bulbar, recto-prostatic and recto-vesical fistulas [15]. One factor that predisposes occurrence of urinary calculi in patients with ARM is incomplete fistula resection with posterior diverticulum lithiasis [15]. Additionally, the communication of the urinary tract to the rectum in patients with such fistulae predisposes the urinary tract to infections which may form a nidus to bladder stone formation [16].

We acknowledge that data was collected from patient records, consequently any errors of documentation during initial recording of these details contribute to our study limitations. Additionally, the study population is small and therefore not representative of a true epidemiological study. To this effect the study herein may not make conclusive statements on epidemiological statistics rather the study has made inferences that can be tested in larger study populations.

Conclusion

This study reveals a general inclination towards delayed diagnosis in children presenting with bladder stones especially in the absence of acute urinary retention and hematuria. Additionally, congenital anomalies of the urinary tract should be suspected in patients with bladder stones particularly in presence of urinary tract dilatation. Finally, ARM presents a unique and rare association with bladder calculi in children.

Acknowledgment

Kenyatta National Hospital, Department of Specialized surgery, Paediatric Surgery Unit.

University of Nairobi, School of Medicine, Thematic Unit of Paediatric Surgery.

Funding Support

No funding or grant support.

Conflict of Interest

The following authors have no financial disclosures: T.J, J.L.

Bibliography

1. Lal Bhamar, *et al.* "Childhood Bladder Stones-An Endemic Disease of Developing Countries". *Journal of Ayub Medical College, Abbottabad: JAMC* 27.1 (2015): 17-21.

2. Halstead Scott B. "Epidemiology of bladder stone of children: precipitating events". *Urolithiasis* 44.2 (2016): 101-108.
3. Philippou Prodromos., *et al.* "The management of bladder lithiasis in the modern era of endourology". *Urology* 79.5 (2012): 980-986.
4. Donaldson James F., *et al.* "Treatment of Bladder Stones in Adults and Children: A Systematic Review and Meta-analysis on Behalf of the European Association of Urology Urolithiasis Guideline Panel". *European Urology* 76.3 (2019): 352-367.
5. Soliman Neveen A and S Adibul Hasan Rizvi. "Endemic bladder calculi in children". *Pediatric Nephrology* 32.9 (2017): 1489-1499.
6. Ngugi PM., *et al.* "Urolithiasis in Nairobi, Kenya". *East African Medical Journal* 87.10 (2010): 395-399.
7. Njoka JT., *et al.* "Kenya: Country situation Assessment. Working paper. Pathways to Resilience in Semi-arid Economies (PRISE) project (2016).
8. Al-Marhoon Mohammed S., *et al.* "Comparison of endourological and open cystolithotomy in the management of bladder stones in children". *The Journal of Urology* 181.6 (2009): 2684-2687.
9. Wambui Winfred Muringi., *et al.* "Determinants of Health Seeking Behavior among Caregivers of Infants Admitted with Acute Childhood Illnesses at Kenyatta National Hospital, Nairobi, Kenya". *International Journal of Pediatrics* (2018): 5190287.
10. Madani Abbas., *et al.* "Urinary calcium and uric acid excretion in children with vesicoureteral reflux". *Pediatric Nephrology* 27.1 (2012): 95-99.
11. Mahyar Abolfazl., *et al.* "The association of hypercalciuria and hyperuricosuria with vesicoureteral reflux in children". *Clinical and Experimental Nephrology* 21.1 (2017): 112-116.
12. Adedoyin OT., *et al.* "Bladder stone in a three-year-old Nigerian child with posterior urethral valve: a case report". *West African Journal of Medicine* 30.3 (2011): 214-216.
13. Acharya Himanshu., *et al.* "Posterior urethral valves with vesical calculus: A rare association". *Journal of Indian Association of Pediatric Surgeons* 14.3 (2009): 115-116.
14. Méndez-Gallart Roberto., *et al.* "Urinary retention in an infant caused by a posterior urethral diverticulum lithiasis following surgery for anorectal malformation". *Urolithiasis* 42.2 (2014): 183-185.
15. Levitt Marc A and Alberto Peña. "Anorectal malformations". *Orphanet Journal of Rare Diseases* 2.33 (2007).
16. Schwaderer Andrew L and Alan J Wolfe. "The association between bacteria and urinary stones". *Annals of Translational Medicine* 5.2 (2017): 32.

Volume 4 Issue 11 November 2021

© All rights are reserved by Jumbi T and Lessan J.