

Determination of Some Elements in Gall Stones and Some Natural Samples (Tap Water, Black Tea, and Table Salt (Na Cl))

Hiba HT, Mohammed SA, Ishraga EA and Abdal Salam AD*

Abdal Salam AD, Department of Chemistry, Faculty of Science, Sudan University of Science and Technology, Khartoum, Sudan

*Corresponding Author: Abdal Salam AD, Department of Chemistry, Faculty of Science, Sudan University of Science and Technology, Khartoum, Sudan.

Received: September 26, 2019; Published: November 04, 2019

DOI: 10.31080/ASPS.2019.03.0431

Abstract

The high amount or decreases in the metal ions consumed in the human body can cause a serious problem on human health. For this problem inductively coupled plasma optical emission (ICP) was used to analyze human gallstones elements (Ba^{2+} , Ca^{2+} , Cu^{2+} , Fe^{2+} , K^{1+} , Mg^{2+} , Mn^{2+} , Na^{1+} and Se^{2+}) and some natural samples (tap water, black tea and sodium chloride (Na Cl)). Ten gall stones were collected during surgery from patients with cholesterol, pigment and mixed stones. The results of quantitative elemental analysis revealed that, Ca^{2+} was high in samples 4,5,7,8 and black tea (12), K^{1+} was high in black tea (12) and Sodium chloride (13), Mg^{2+} was high in black tea (12) and Sodium chloride (13), Na^{1+} was high in Sodium chloride (13). The rate of elements (Ca^{2+} , K^{1+} , Na^{1+} , Mg^{2+}) were high in black tea and sodium chloride probably that reason indicates the stone formation in the human body.

Keywords: Inductively Coupled Plasma Mass Spectrometry (ICP-MS); Gallstones; Cholesterol Stones; Pigment Stones; Mixed Stones

Introduction

Gallstones are frequently composed of more than one crystalline compound, although the exact mechanism of formation of multicomponent stones is not clearly understood. They are generally classified into three major types: pure cholesterol, pigment and mixed gallstone. Pigment gallstones are further subdivided into laminated brown stones and amorphous black stones [1]. Cholesterol gallstones are higher prevalence than pigment gallstones. Diet affected the type of gallstone formation. Studies reported that Consumption of fried foods and animal lipid increased risk of cholesterol gallstone formation, while intake of carbohydrate increases risk of pigment gallstones formation. Suggesting diet influenced the type of gallstone formation [2]. Aetiology and pathogenesis of cholesterol gallstones still are not well defined, and strategies for prevention and efficient nonsurgical therapies are missing. This review summarizes current concepts on the pathogenesis of cholesterol gallstones with focus on the uptake and secretion of biliary lipids and special emphasis on recent studies into the genetic background. More than 80% of gallstone carriers are unaware of their gallbladder disease [3,4]. Gallstone disease is the most common gastrointestinal disorder affecting the biliary system. Its complications are caused by inflammation, infection, or ductal obstruction. Gallbladder stone disease is one of the major surgical problems in several populations and it accounts for many hospital admissions

and surgical interventions. Gallstones represent a major problem in many countries [5]. Deposition of hard fatty or mineral in the gallbladder leads to Gallstone disease. In this disease, the stones block part of biliary system and cause irritation of gallbladder and complications. This disorder can cause life threatening conditions if their symptoms or complications left without treatment [6]. About 1–2% per year of patients develop complications and need surgery [7]. The prevalence of cholelithiasis in Asian population is estimated to be 3% to 10%. Published data showed the prevalence of GSD to be 3.2% in Japan [8], 3~11% in China [9], 7.1% in Northern India [10] and 5% in Taiwan [11]. In Pakistan, gallstones are found in 8% and 20% of patients above 40 and 60 years respectively [12]. In the US, gallstone disease has the most common inpatient diagnosis among gastrointestinal and liver diseases [8] and stands for \$5.8 billion direct costs, exceeded only by gastro esophageal reflux disease [13]. Cholesterol stones contain more than 70% of cholesterol whereas pigment stones contain mainly of various bilirubinate salts with less than 20% cholesterol by weight. Pigment stones are predominant in India [14]. In this study we determine some elements in gall stones and some natural samples (tap water, black tea, and table salt (Na Cl)) we found that the reason of the stone formation in the human body from black tea.

Materials and Method

- Samples: Gall stones were collected from 10 Sudanese patients in Sudan. The stones were collected, dried and powdered to homogenous mixture and transferred into plastic containers.
- Chemicals: All chemicals used in this work were of analytical grade type (AR), Hydrochloric acid (36-38%), Nitric acid (96-72%), anhydrous potassium bromide, de ionized water.
- Equipment: Mortar, Beakers, volumetric flask.
- Instruments: Varian 725-ES inductively coupled plasma optical emission spectrometer.
- Methods: (0.5 g) of the collected gall stones was placed into digestive vial and 9ml of aquilegia (6ml HCl conc. and 2 mL

HNO3 conc.) was added. The volume was then adjusted to 25 ml by adding water and the mixture was shaken. Na cl sample solution was prepared by dissolving 10mg of powdered sample in demonized water and diluted to 100 ml with demonized water in volumetric flask. Black tea sample solution was prepared by boiling 10mg of sample in demonized water; it was filtrated and diluted to 100 ml with demonized water in volumetric flask. All samples were analyzed by inductively coupled plasma optical emission (ICP), varin, and 725-ES.

Results

The elements content of different gall stones samples tap water, black tea and sodium chloride were as shown in table 1 given by Inductively Coupled Plasma- Optical Emission Spectrometer (ICP), varin, 725-ES.

Samples	Age	Ba ²⁺	Ca ²⁺	Cu ²⁺	Fe ²⁺	K ¹⁺	Mg ²⁺	Mn ²⁺	Na ¹⁺	Se ²⁺
Samples 1	45	43.54	52.660	35.69	20.93	0.0318	354.0	14.3.7	151.2	0.0266
Samples 2	40	0.0005	29.310	545.6	54.61	0.0318	247.7	149.6	254.3	0.0266
Samples 3	54	1.569	1.371	8.776	96.75	0.031	45.37	4.117	19.57	0.0266
Samples 4	26	1.458	732	2.893	37.14	163.8	53.11	0.3205	48.99	0.0266
Samples 5	68	2.881	68.620	4172	1418	399	1021	424.3	277.6	0.0266
Samples 6	53	0.0005	15.990	723.0	189.29	458.0	631.5	139.5	269.0	0.0266
Samples 7	45	0.0005	88.980	464.3	280.5	47.64	927.7	1131	234.3	0.0266
Samples 8	44	0.0005	201.000	151.9	145.7	82.38	837.4	121.1	172.1	0.0266
Samples 9	20	0.0005	17.710	483.8	175.8	155.8	198.7	57.4	355.6	0.0266
Samples 10	50	0.0005	141.300	1438	325.1	386.3	2337	3025	449.1	119.1
Samples 11 Tap water		0.1813	54.66	0	0.0344	5.326	18.05	0	32.38	0
Samples 12 Black tea		8.7847	2192	34.35	40.24	17042	1207	648.0	700	12.37
Samples 13 Na Cl		0.9478	0.9478	0	0	740	1743	3.227	355000	0

Table 1: As we can see, the content average level (in p.p.m).

Figure 1: Ba²⁺ Content (p.p.m) by ICP in sample1-13.

Figure 2: Ca²⁺ Content (p.p.m) by ICP in sample1-13..

Figure 3: Cu²⁺ Content (p.p.m) by ICP in sample1-13.

Figure 7: Mn²⁺ Content (p.p.m) by ICP in sample1-13.

Figure 4: Fe²⁺ Content (p.p.m) by ICP in sample1-13.

Figure 8: Na¹⁺ Content (p.p.m) by ICP in sample1-1

Figure 5: K¹⁺ Content (p.p.m) by ICP in sample1-13.

Figure 9: Se²⁺ Content (p.p.m) by ICP in sample1-13.

Figure 6: Mg²⁺ Content (p.p.m) by ICP in sample1-13.

Discussion

The result of elements of the different gall stones samples (1-10), tap water (1-1), black tea (12), and sodium chloride (Na Cl) (13), gave indication to the concentration of element, the rate of (Ba) was high in samples (1,3,4,5, and black tea (12), the rate of (Ca) was high in samples (4,5,7,8 and black tea (12), the rate of (Cu) was high in samples (2,5,6,9 and 10, the rate of (Fe) was high in samples (5,6,7,9 and 10). The rate of (K) was high in ample 13 and black tea (12), the rate of (Mg) was high in sample (10, black

tea (12), and sodium chloride (13)) the rate of (Mn) was high in samples (7,10, and black tea (12). the rate of Na was high in sodium chloride (13), the rate of (Se) was high in samples (10 and in the black tea (12). The rate of elements (Ca, K, Na, Mg,) were high in black tea and sodium chloride probably that reason indicates the stone formation. The gallbladder can develop only one large stone or hundreds of small stones, or a combination of the two. Gallstones formation in not related to single reason. Cholesterol stones formed when the liver produces much cholesterol in the bile which form crystals and changed to hard stones in the gallbladder. Pigment stones they differ in their chemical composition and colored (black and brown) by the presence of bilirubin, the pigment in RBCs. They are composed of bilirubin polymers or calcium salts. These stones formed because of changes in bile components or due to inability of gallbladder to empty normally. Pigment stones are associated with chronic hemolysis in conditions such as sickle cell disease, thalassemia, and cirrhosis. Mixed stones composed of cholesterol and salts. These types are the most common. Gallstones formation become more common with older age, female gender, overweight and diabetic people [15,16].

Conclusion

The result of the elements concentration of different gall stone samples (1-10), tap water (11), black tea, (12) and sodium chloride (13) gave indication to the concentration of element, the rate of element was high in black tea sample, probably, that was the reason indicate that the stone formation.

Bibliography

1. Beckingham IJ. "ABC of diseases of liver, pancreas, and biliary system. Gallstone disease". *BMJ* 322 (2001): 91-94.
2. Bashir M Jarrar and Meshref A Al-Rowaili. "Chemical Composition of Gallstones from Al-Jouf Province of Saudi Arabia". *Malaysian Journal of Medical Sciences* 18 (2011): 47-52.
3. UAttili AF, *et al.* "Epidemiology of gallstone disease in Italy: prevalence data of the Multicenter Italian Study on Cholelithiasis (M.I.COL)". *American Journal of Epidemiology* 141 (1995): 158-165.
4. Heaton KW, *et al.* "Symptomatic and silent gall stones in the community". *Gut* 32 (1991): 316-320.
5. Yomna Ali, *et al.* "Gallbladder Stone Disease and its Relation to Dietary Intake" (2019).
6. Li Huang, *et al.* "Farnesoid X-receptor activates transcription of the phospholipid pump MDR3". *Journal of Biological Chemistry* 278 (2003): 51085-51090.
7. Friedman GD, *et al.* "Prognosis of gallstones with mild or no symptoms: 25 years of follow-up in a health maintenance organization". *Journal of Clinical Epidemiology* 42 (1989): 127-136.
8. Nomura H, *et al.* "Prevalence of gallstone disease in a general population of Okinawa, Japan". *American Journal of Epidemiology* 128 (1988): 598-605.
9. Zhang W, *et al.* "Epidemiology and risk factors of cholelithiasis". *Journal of Surgery Concepts and Practice* 16 (2011): 408-412.
10. Unisa S, *et al.* "Population-based study to estimate the prevalence and determine risk factors of gallbladder diseases in the rural Gangetic of North India". *HPB (Oxford)* 13 (2011): 117-125.
11. Chen CH, *et al.* "Prevalence and risk factors of gallstone disease in an adult population of Taiwan: an epidemiological survey". *Journal of Gastroenterology and Hepatology* 21 (2006): 1737-1743.
12. Moerman CJ, *et al.* "Dietary risk factors for clinically diagnosed gallstones in middle-aged men. A 25-year follow-up study (the Zutphen Study)". *Annals of Epidemiology* 4 (1994): 248-254.
13. Sandler RS, *et al.* "The burden of selected digestive diseases in the United States". *Gastroenterology* 122 (2002): 1500-1511.
14. Jayanthi V, *et al.* "Epidemiology of gallstone disease –top line findings". *Bombay hospital Journal* 41 (1999) 495-502.
15. Saad Muhmood Hussain, *et al.* "Determination of Chemical Composition of Gallbladder Stones and their Association with Induction of Cholangiocarcinoma". *Asian Pacific Journal of Cancer Prevention* 14 (2013): 6257-6260.
16. Mohammed Helmy Faris Shalayel, *et al.* "Biochemical composition of gallstones: Do different genders differ". *American Journal of Biological Chemistry* 1 (2013): 1-6.

Volume 3 Issue 12 December 2019

© All rights are reserved by Abdal Salam AD., *et al.*